

**PrEP**

# *Daily PrEP to prevent HIV*

**If you are HIV-negative and concerned about exposure to HIV, ask your doctor if PrEP (Pre-exposure Prophylaxis) may be right for you.**

PrEP is a daily pill that can protect you from HIV as long as you take it every day.

Condoms provide additional protection against HIV, most sexually transmitted infections and unintended pregnancy.

Many insurance plans, including Medicaid, cover PrEP. Assistance may be available if you are uninsured.

**Call 311 or visit [nyc.gov](http://nyc.gov) and search “HIV PrEP and PEP” for more information.**





PEP

## *Emergency PEP to prevent HIV*

If you are HIV-negative and think you were exposed to HIV, **immediately** go to a clinic or emergency room and ask for **PEP** (Post-exposure Prophylaxis).

PEP is an emergency medicine you take right after you are exposed to HIV.

PEP can stop HIV if started within 36 hours of exposure. You continue taking PEP for 28 days.

Many insurance plans, including Medicaid, cover PEP. Assistance may be available if you are uninsured.

Call 311 or visit [nyc.gov](http://nyc.gov) and search “HIV PrEP and PEP” for more information.



**NYC**  
Health



**PrEP**

# *Daily PrEP to prevent HIV*

**If you are HIV-negative and concerned about exposure to HIV, ask your doctor if PrEP (Pre-exposure Prophylaxis) may be right for you.**

PrEP is a daily pill that can protect you from HIV as long as you take it every day.

Condoms provide additional protection against HIV, most sexually transmitted infections and unintended pregnancy.

Many insurance plans, including Medicaid, cover PrEP. Assistance may be available if you are uninsured.

**Call 311 or visit [nyc.gov](http://nyc.gov) and search “HIV PrEP and PEP” for more information.**



**NYC**  
Health



PEP

## *Emergency PEP to prevent HIV*

If you are HIV-negative and think you were exposed to HIV, **immediately go** to a clinic or emergency room and ask for **PEP** (Post-exposure Prophylaxis).

PEP is an emergency medicine you take right after you are exposed to HIV.

PEP can stop HIV if started within 36 hours of exposure. You continue taking PEP for 28 days.

Many insurance plans, including Medicaid, cover PEP. Assistance may be available if you are uninsured.

Call 311 or visit [nyc.gov](http://nyc.gov) and search “HIV PrEP and PEP” for more information.



**NYC**  
Health



**PrEP**

# *Daily PrEP to prevent HIV*

**If you are HIV-negative and concerned about exposure to HIV, ask your doctor if PrEP (Pre-exposure Prophylaxis) may be right for you.**

PrEP is a daily pill that can protect you from HIV as long as you take it every day.

Condoms provide additional protection against HIV, most sexually transmitted infections and unintended pregnancy.

Many insurance plans, including Medicaid, cover PrEP. Assistance may be available if you are uninsured.

**Call 311 or visit [nyc.gov](http://nyc.gov) and search “HIV PrEP and PEP” for more information.**



**NYC**  
Health



PEP

## *Emergency PEP to prevent HIV*

If you are HIV-negative and think you were exposed to HIV, **immediately go** to a clinic or emergency room and ask for **PEP** (Post-exposure Prophylaxis).

PEP is an emergency medicine you take right after you are exposed to HIV.

PEP can stop HIV if started within 36 hours of exposure. You continue taking PEP for 28 days.

Many insurance plans, including Medicaid, cover PEP. Assistance may be available if you are uninsured.

Call 311 or visit [nyc.gov](http://nyc.gov) and search “HIV PrEP and PEP” for more information.



**NYC**  
Health