



**NEW YORK CITY DEPARTMENT OF  
HEALTH AND MENTAL HYGIENE**

Mary T. Bassett, MD, MPH  
Commissioner

Joseph Burzynski, MD, MPH  
Assistant Commissioner  
[JBurzyns@health.nyc.gov](mailto:JBurzyns@health.nyc.gov)

Bureau of Tuberculosis Control  
42-09 28th Street, CN-72B,  
Long Island City, NY 11101

October 24, 2017

To Whom It May Concern:

A person whose test result for tuberculosis (TB) infection is positive should undergo a chest x-ray to rule out active TB disease. A physician uses the chest x-ray result, as well as the medical history and physical examination, to determine appropriate follow up.

**Once a person:**

- **tests positive for TB infection and**
- **has had a chest x-ray that is negative for active TB disease**

**no follow-up radiologic tests are recommended or indicated as long as the person has no symptoms of active TB.**

Some employers still recommend routine follow-up chest x-rays at regular intervals; such radiographs are not indicated, nor are they recommended by city, state or federal health agencies.<sup>1,2</sup>

The New York City Department of Health and Mental Hygiene Chest Centers will not perform follow-up chest x-rays for persons who are already known to have a positive test for TB infection, have had a chest x-ray that is negative for active TB disease, and who have no symptoms of active TB. If completion of a follow-up chest x-ray is a condition of employment, individuals or employers may contact the Bureau of Tuberculosis Control (BTBC) on the TB Hotline at (844) 713-0559 for assistance to bring their policies into accordance with city, state and federal recommendations.

Chest x-rays should be obtained on all patients with suspected or confirmed active TB as indicated, regardless of when the test for TB infection was done. In addition, a chest X-ray should be performed just prior to the initiation of prophylaxis for TB infection.

Sincerely,

Joseph Burzynski, MD, MPH  
Assistant Commissioner  
Bureau of Tuberculosis Control

1 New York City Department of Health and Mental Hygiene, Bureau of Tuberculosis Control. *Clinical Policies and Protocols*, 3rd ed., 1999.

2 Centers for Disease Control and Prevention. Guidelines for preventing the transmission of *Mycobacterium tuberculosis* in Health Care Facilities, 1994. *MMWR* 1994;43(RR-3):40.