Source: New York City, Office of	Chief Medical Examiner (OC 2013		
Variables	N	2013	AAR
Total	788	100%	11.6
Gender			
Male	570	72%	17.7
Female	218	28%	6.2
Race/Ethnicity ²			
Black	172	22%	10.7
Hispanic	222	28%	12.1
White Other	358 36	45% 5%	16.3 X
	30	5%	~
Age-group ³	45	C 0/	4.0
15-24 25-34	45 137	6% 17%	4.0 9.3
25-34 35-44	157	20%	9.3 13.6
45-54	251	32%	22.6
43-34 55-64	163	21%	17.1
65-84	33	4%	3.6
Unknown	00	770	0.0
Mean Age	45.3		
Borough of Residence ⁴			
Bronx	162	21%	15.0
Brooklyn	165	21%	8.2
Manhattan	136	17%	9.8
Queens	145	18%	7.6
SI	64	8%	17.6
Other	116	15%	Х
Borough of Death			
Bronx	184	23%	17.2
Brooklyn	197	25%	9.9
Manhattan	191	24%	13.8
Queens	152	19%	8.0
SI	64	8%	17.5
Neighborhood poverty ⁵			
Group 1: 0 to <10%	147	19%	10.6
Group 2: 10 to <20%	183	23%	7.1
Group 3: 20 to <30%	160	20%	10.0
Group 4: 30 to 100%	181	23%	15.9
Unknown/Other	117	15%	Х

 Table 1. Demographics of unintentional drug poisoning decedents, age-adjusted rates¹ per 100,000 residents, New York City, 2013

 Source: New York City, Office of Chief Medical Examiner (OCME)

¹Rates are calculated using intercensal New York City population denominators updated December 2014. Rates are adjusted to 2000 US Standard population.

²Other race/ethnicity or missing race/ethnicity are collapsed into "Other", and are not included in the percent of total calculation.

³Age standardized rates are presented. Unknown age are not included in the percent of total calculation.

⁴Decedents with missing residence information or non-NYC residence on death certificate are collapsed into "Other" and are not included in the percent of total calculation.

⁵Neighborhood poverty (based on UHF) defined as percent of residents with incomes below 100% of the Federal Poverty Level per American Community Survey Census 2000.

