Table 1. Demographic and clinical characteristics of unique patients with alcohol-related hospitalizations¹, age adjusted rates (AAR)² per 100,000 residents, New York City, 2011 Source: New York State Department of Health, Statewide Planning and Research Cooperative System (SPARCS), 2011 (Data update: April 2014).

| Course: Now York Clare Departm | 2011 | | |
|--|------------------|--------|----------------|
| | Unique Patients | | |
| | N | % | AAR |
| Total | 41,229 | 67% | 598.5 |
| Type of diagnosis ³ | 41,223 | 01 /0 | 330.3 |
| Dependence | 17,294 | 42% | 251.5 |
| Abuse | 19,800 | 48% | 287.6 |
| Alcohol induced mental disorder | 6,766 | 16% | 98.7 |
| Alcohol liver and cirrhosis | 4,713 | 11% | 67.9 |
| Alcohol other | 9,937 | 24% | 143.1 |
| Gender | 3,337 | 2470 | 140.1 |
| Female | 12,154 | 29% | 329.4 |
| Male | 29,074 | 71% | 913.4 |
| Race/Ethnicity ⁴ | 23,014 | 7 1 70 | 313.4 |
| Black | 13,619 | 33% | 850.4 |
| Hispanic | 8,790 | 21% | 500.9 |
| White | 10,016 | 24% | 406.0 |
| Other | 8,804 | 21% | 858.6 |
| | 0,004 | 2170 | 030.0 |
| Age-group ⁵ 13-17 | 380 | 1% | 78.7 |
| 18-24 | 1,762 | 4% | 204.6 |
| 25-34 | 4,371 | 11% | 306.0 |
| 35-44 | 6,153 | 15% | 531.3 |
| 45-54 | 10,981 | 27% | 985.8 |
| 55-64 | 9,791 | 24% | 1060.2 |
| 65-84 | 7,791 | 19% | 896.0 |
| Borough of Residence | 7,791 | 1970 | 090.0 |
| Bronx | 10.027 | 24% | 020.2 |
| | 10,037 12,724 | 31% | 928.2 623.5 |
| Brooklyn Manhattan | 8,563 | 21% | 617.4 |
| Queens | 7,640 | 19% | 393.7 |
| Staten Island | · · | 19% | 552.0 |
| | 2,265 | 3% | 552.0 |
| Borough of Hospital | 0.000 | 220/ | 004.0 |
| Bronx | 8,909 | 22% | 824.6 |
| Brooklyn | 11,329 | 27% | 555.4 |
| Manhattan | 12,581 | 31% | 907.9 |
| Queens | 6,358 | 15% | 327.7 |
| Staten Island | 2,052 | 5% | 501.2 |
| Neighborhood poverty ⁶ | 2 224 | 001 | 202.2 |
| Group 1: 0 to <10% | 3,334 | 8% | 399.3 |
| Group 2: 10 to <20% | 16,017 | 39% | 462.5 |
| Group 3: 20 to <30% | 11,066 | 27% | 691.2 |
| Group 4: 30 to 100% Excludes detox and rehabilitation (94.6), suice | 10,729 | 26% | 1109.2 |

¹ Excludes detox and rehabilitation (94.6), suicide (E950), attempted assault (E960), and poisoning by solid or liquid substances undetermined whether accidentally or purposely inflicted (E980).



²Rates are calculated using intercensal New York City population denominators updated April 2014. Rates are adjusted to 2000 US Standard population.

³Ever diagnosed in a given year. Diagnoses are not mutually exclusive.

⁴Other race/ethnicity or missing race/ethnicity are collapsed into "Other", and are not included in the percent of total calculation.

⁵Age standardized rates are presented. Unknown age are not included in the percent of total calculation.

⁶Neighborhood poverty (based on UHF) defined as percent of residents with incomes below 100% of the Federal Poverty Level per American Community Survey Census 2009-2012.