

## DETECTING AND TREATING DEPRESSION IN ADULTS

### REFERENCES

1. Ballenger JC, Davidson JR, Lecrubier Y, et al. Consensus statement on the primary care management of depression from the International Consensus Group on Depression and Anxiety. *J Clin Psychiatry*. 1999;60(suppl 7):54-61.
2. Robins LN, Helzer JE, Weissman MM, et al. Lifetime prevalence of specific psychiatric disorders in three sites. *Arch Gen Psychiatry*. 1984;41:949-958.
3. Johnson J, Weissman MM, Klerman GL. Service utilization and social morbidity associated with depressive symptoms in the community. *JAMA*. 1992;267:1478-1483.
4. Horowitz JA, Goodman JH. Identifying and treating postpartum depression. *J Obstet Gynecol Neonatal Nurs* 2005;(2):264-273.
5. Katon W, Shulberg H. Epidemiology of depression in primary care. *General Hospital Psychiatry*. 1992; 14:237-242.
6. Conwell Y. Suicide in elderly patients. In: Schneider LS, Reynolds CF III, Lebowitz BD, Friedhoff A, eds. *Diagnosis and treatment of depression in late life: Results of the NIH Consensus Development Conference*. Washington, DC: American Psychiatric Publishing; 1994:397-418.
7. Arroll B, Khin N, Kerse N. Screening for depression in primary care with two verbally asked questions: cross sectional study. *BMJ*. 2003;327:1144-1146.
8. Simon GE, VonKorff M, Piccinelli M, Fullerton C, Ormel J. An international study of the relation between somatic symptoms and depression *N Engl J Med*. 1999;341(18):1329-1335.
9. Kroenke K, Spitzer RL, Williams JB. The PHQ-9. Validity of a brief depression severity measure. *J Gen Intern Med*. 2001;16:606-613.
10. *Diagnostic and Statistical Manual of Mental Disorders*. 4th ed. text rev. Washington, DC: American Psychiatric Association; 2000.
11. Practice guideline for the assessment and treatment of patients with suicidal behaviors. *Am J Psychiatry*. 2003;160(suppl 11):1-60.
12. Intermountain Health Care. Depression 2004 Update. Available at [www.ihc.com/xp/ihc/documents/clinical/103/8/5/cpmdepression.pdf](http://www.ihc.com/xp/ihc/documents/clinical/103/8/5/cpmdepression.pdf). Accessed October 21, 2005.
13. Agency for Health Care Policy and Research. Clinical Practice Guideline No. 5: Depression in Primary Care: Detection, Diagnosis and Treatment. Rockville, Md: Public Health Service, U.S. Dept of Health and Human Services; 1993. AHCPR publication 93-0550.
14. Brown GK, Bruce ML, Pearson JL. High-risk management guidelines for elderly suicidal patients in primary care settings. *Int J Geriatr Psychiatry*. 2001;16:593-601.
15. Stuart MR, Lieberman JA. *The Fifteen Minute Hour: Applied Psychotherapy for the Primary Care Physician*. 2nd ed. Westport, Conn: Greenwood Publishing Group; 1993.
16. Duloxetine (Cymbalta): A new SNRI for depression. *The Medical Letter*. October 2004;46(1193):81-83.
17. Treatment Guidelines from the Medical Letter. Drugs for psychiatric disorders. *The Medical Letter*. July 2003; 11:69-76.
18. American Psychiatric Association. Practice guideline for the treatment of patients with bipolar disorder. Available at [www.psych.org/psych\\_pract/treatg/pg/Practice%20Guidelines8904/MajorDepressiveDisorder\\_2e.pdf](http://www.psych.org/psych_pract/treatg/pg/Practice%20Guidelines8904/MajorDepressiveDisorder_2e.pdf). Accessed November 2, 2005.