


Community Health Profiles

New York City Department of Health and Mental Hygiene

SECOND EDITION — 2006



TAKE CARE Northeast Queens

*(Including Bayside,
Douglaston, Little Neck,
and Oakland Gardens)*

Community Health Profile, Second Edition: Northeast Queens

New York City is the most diverse city in the U.S. — a fact reflected in the distinct character of each neighborhood. The second edition of the Community Health Profiles uses **Take Care New York** (TCNY), the city’s health policy, to examine preventable causes of illness and death in all of NYC’s 42 neighborhoods. This report updates the 2002 profile (available at nyc.gov/health) by providing more recent and time-trend data, and a greater variety of health statistics. Key health issues in **Northeast Queens** include:

- Cancer is the primary cause of premature death in Northeast Queens (page 4).
- Cancer screening rates among women and older adults in Northeast Queens remain below TCNY targets (page 11).
- Hospitalizations for fall-related injuries — particularly hip fractures — among older adults are more common in Northeast Queens than in New York City overall (page 14).

Methods: While this report provides important information, it is not intended to be an exhaustive examination of the health of Northeast Queens residents, as not all health problems and their causes could be covered. Only statistically significant findings are discussed in the text. For complete information on methods, see Technical Notes (page 15).

Northeast Queens at a Glance

Population

Total number of people living in Northeast Queens in 2000:

88,200


Age

People in Northeast Queens are slightly older than in Queens and New York City overall

	Northeast Queens	Queens	NYC
0-17 years	20%	23%	24%
18-24 years	7%	9%	10%
25-44 years	30%	33%	33%
45-64 years	26%	22%	21%
65+ years	17%	13%	12%

Poverty

In Northeast Queens, the percent of residents living below the poverty level is lower than in Queens and NYC overall



Education

Northeast Queens residents aged 25 and older have completed more years of education than those in NYC overall

	Northeast Queens	Queens	NYC
Up to 8th grade	6%	11%	12%
Some high school, no diploma	8%	15%	16%
High school diploma	24%	28%	25%
Some college, no degree	23%	22%	20%
College graduate	39%	24%	27%




Foreign-born

More than one third of Northeast Queens residents were born outside the U.S.

Area	Percent
Northeast Queens	35%
Queens	46%
NYC	36%

Race / Ethnicity

Northeast Queens has higher proportions of white and Asian residents than Queens and NYC overall



Data Source: U.S. Census 2000/NYC Department of City Planning

Take Care Northeast Queens



In 2004, the Health Department created a citywide health policy called **Take Care New York** (TCNY) to help improve the health of New Yorkers. TCNY identifies 10 key areas that cause significant illness and death but can be improved through intervention by individuals, health care providers, government agencies, and other organizations.

This report examines how well Northeast Queens residents are doing on health indicators for each of the 10 TCNY goals. It examines areas in which the community is a health leader, as well as areas that need improvement. The TCNY report card below shows where Northeast Queens ranks among all 42 New York City neighborhoods. (See Technical Notes for information about how neighborhoods were defined and ranked.)

Take Care New York report card


Northeast Queens ranks as average or above on all indicators when compared to the 41 other NYC neighborhoods

	Below Average (bottom 10)	Average (middle 22)	Above Average (top 10)
Take Care New York Goals			
1 Have a regular doctor			✓
2 Be tobacco-free		✓	
3 Keep your heart healthy			✓
4 Know your HIV status		✓	
5 Get help for depression			✓
6 Live free of alcohol and drugs			✓
7 Get checked for cancer			✓
8 Get the immunizations you need		✓	
9 Make your home safe and healthy			✓
10 Have a healthy baby			✓

How Residents Rate Their Own Health

Overall health

Adults in Northeast Queens are less likely to consider themselves to be in fair or poor health



Percents are age-adjusted.
Data Source: NYC Community Health Survey 2002-03-04


People are good at rating their own health. In general, when asked to rate their general health as excellent, very good, good, fair, or poor, those who say “fair” or “poor” are more likely to have health problems than those who report better health.

In Northeast Queens, residents are less likely to report being in fair or poor health (13%) than those both in Queens (20%) and in New York City overall (21%).

Overall Death Rates in Northeast Queens

Death rates

In Northeast Queens, death rates are lower than in Queens and NYC overall



Rates are age-adjusted.
Data Sources: Bureau of Vital Statistics, NYC DOHMH, 1995-2004; U.S. Census 1990 and 2000/NYC Department of City Planning

The death rate in Northeast Queens has declined slightly in the past decade, mirroring the rate drop in New York City overall.

In 2003-2004, the average annual death rate in Northeast Queens was 25% lower than in Queens and 35% lower than in New York City overall (461/100,000 vs. 610/100,000 in Queens and 718/100,000 in NYC). Throughout this profile, cause-specific death rates are provided for TCNY goals.

Line graphs. All time-trend data are presented as annual averages with 2 or 3 years of data combined. For example, in this graph, the first point on each line represents the average annual death rate for 1995 and 1996 combined.


Premature death

People who die before age 75 can be thought of as dying early, or prematurely. If a person dies early, their years of potential life lost (YPLLs) can be calculated by subtracting their age at death from 75 years to get a measure of premature death.

The causes of premature death differ across communities. The primary cause of premature death in Northeast Queens is cancer, as well as in both Queens and New York City overall.

Top 5 causes of years of potential life lost

Cancer causes the most years of potential life lost in Northeast Queens




*Other includes Diabetes (3%), Pneumonia and Influenza (2%), Chronic Lower Respiratory Disease (2%), Congenital Conditions (2%), Unknown or Benign Neoplasms (2%), and Other (15%).

Data Source: Bureau of Vital Statistics, NYC DOHMH, 2002-04

Death before age 75

The 2003-2004 average annual death rate for people younger than 75 years in Northeast Queens is the lowest, ranking 1st among 42 NYC neighborhoods



Data Sources: Bureau of Vital Statistics, NYC DOHMH, 2003-04; U.S. Census 2000/NYC Department of City Planning



Take Care New York Goals


GOAL 1 Have a Regular Doctor or Other Health Care Provider

Access to good medical care helps people prevent illnesses, identify health conditions early, and treat health problems. Some conditions can and should be managed regularly outside the hospital. Higher rates of these avoidable hospitalizations can indicate reduced access to health care in a community.

Having a “medical home” — a personal doctor or other health care provider and a regular place of care other than the emergency department (ED) — is a critical component of good health care access. Northeast Queens residents are less likely to be without a regular doctor than adults in both Queens and NYC overall, meeting the TCNY goal of less than 20%. Northeast Queens residents are also less likely to go to the ED when they are sick or need health advice (2%) than those in Queens (7%) and NYC overall (8%).

Access to care


Without a primary provider, people may seek routine health care in the emergency department (ED)



*Estimate is unstable due to small sample size and should be interpreted with caution. Percents are age-adjusted. Data Sources: ¹NYC Community Health Survey 2002-03-04, ²NYC Community Health Survey 2003-04

Avoidable hospitalizations


The 2004 avoidable hospitalization rate in Northeast Queens ranks very well (3rd) among 42 NYC neighborhoods



Data Sources: New York State Department of Health Statewide Planning and Research Cooperative System, 2004; U.S. Census 2000/ NYC Department of City Planning

Health insurance

One in 5 adults in Northeast Queens is uninsured or went without health insurance during the past year



Insurance rates are calculated for adults aged 18-64 and age-adjusted. Data Source: NYC Community Health Survey 2002-03-04

Health insurance is important for access to health care. Northeast Queens adults are more likely to have been insured for the entire past year (80%) than those in Queens (66%) and New York City overall (71%). In addition to the currently uninsured (12%), another 8% of residents in this community went without health insurance at some time during the past year.




GOAL 2 Be Tobacco-Free

Smoking is the leading cause of preventable death in New York City and the cause of many illnesses, including heart disease, stroke, emphysema, and lung cancer. Less than one fifth of Northeast Queens residents (18%) currently smoke. Many methods to quit smoking are available, and 6 in 10 smokers in Northeast Queens (60%) are trying to kick the habit.

Residents who smoke


Fewer than 1 in 5 adults in Northeast Queens smokes



Percents are age-adjusted.
Data Source: NYC Community Health Survey 2004

Attempts to quit smoking in the past year

Most smokers are trying to quit



Percents are age-adjusted.
Data Source: NYC Community Health Survey 2004




GOAL 3 Keep Your Heart Healthy

Heart disease can cause severe illness and death. Northeast Queens residents had an average annual heart disease hospitalization rate in 2003-2004 that was more than 15% lower than the Queens and NYC overall rates (1,458/100,000 vs. 1,727/100,000 in Queens and 1,856/100,000 in NYC). The heart disease death rate in 2003-2004 (221/100,000) was also lower than the rates in Queens (288/100,000) and NYC overall (297/100,000).

Heart disease hospitalizations


Heart disease hospitalizations are less common in Northeast Queens



Rates are age-adjusted.
Data Sources: New York State Department of Health Statewide Planning and Research Cooperative System, 1995-2004; U.S. Census 1990 and 2000/NYC Department of City Planning

Deaths due to heart disease

The heart disease death rate is lower in Northeast Queens




Rates are age-adjusted.
Data Sources: Bureau of Vital Statistics, NYC DOHMH, 1995-2004; U.S. Census 1990 and 2000/NYC Department of City Planning

High blood pressure and high cholesterol. Both of these conditions contribute to heart disease. In Northeast Queens, 26% of adults were told by a health care professional that they have high blood pressure (the same as in Queens and NYC overall), and a quarter (25%) were told that they have high cholesterol (similar to 26% in Queens and NYC overall).

Percents are age-adjusted. Data Source: NYC Community Health Survey 2002

Obesity

More than 1 in 7 adults in Northeast Queens are obese




Obesity is defined as a body-mass-index (BMI) of 30 or greater. Percents are age-adjusted.
Data Source: NYC Community Health Survey 2002-03-04

In addition to smoking, high blood cholesterol and high blood pressure, other factors that put people at risk for heart disease — lack of physical activity and obesity — can be prevented or controlled.

Obesity can lead to a variety of health problems, including heart disease and diabetes. Rates of obesity are increasing rapidly in New York City and across the U.S., making it a major public health concern. In Northeast Queens, 15% of adults are obese.

Diabetes

More than 1 in 20 adults in Northeast Queens have diabetes




Percents are age-adjusted.
Data Source: NYC Community Health Survey 2002-03-04

The increasing prevalence of obesity in the U.S. has contributed to an epidemic of diabetes. About 95% of diabetes cases are type 2 diabetes, which is strongly associated with obesity. Uncontrolled diabetes can worsen the harmful effects of high blood pressure, high cholesterol, and other risk factors for heart disease.

In Northeast Queens, 6% of adults have diabetes.

Days per week of recreational exercise

Only 4 in 10 Northeast Queens adults are meeting physical activity recommendations



Survey Question: On average, how many days per week do you exercise for at least 30 minutes? Percents are age-adjusted.
Data Source: NYC Community Health Survey 2003

Physical activity helps people maintain a healthy weight and strengthens the cardiovascular system. More than 4 in 10 Northeast Queens residents (41%) report doing no physical activity at all. Less than half of residents in this community (40%) report exercising at least 3 days a week.

Centers for Disease Control and Prevention Recommendations

Adults should do either 20 minutes of vigorous exercise 3 times per week or 30 minutes of moderate exercise 5 times per week.




GOAL 4 Know Your HIV Status

Wide disparities exist in HIV across New York City communities. In Northeast Queens, the rate of HIV diagnoses (10/100,000) and the rate of people living with HIV/AIDS (165/100,000) are much lower than the rates in both Queens and NYC overall.

In 2003-2004, the average annual death rate related to HIV disease in Northeast Queens was one sixth of the Queens rate and 95% lower than the NYC overall rate (1/100,000 vs. 6/100,000 in Queens and 18/100,000 in NYC).

Death rate due to HIV

HIV-related death rates in Northeast Queens are much lower than in Queens and NYC overall



HIV/AIDS in 2004

Total HIV diagnoses per 100,000 people* (13+)

Northeast Queens	10
Queens	30
New York City	55

% HIV diagnosed concurrently with AIDS** (13+)

Northeast Queens	50%
Queens	35%
New York City	29%

People living with HIV/AIDS per 100,000 people* (13+)

Northeast Queens	165
Queens	680
New York City	1,419

*Rates are age-adjusted.
**Within 31 days of HIV diagnosis – crude percents
Data Source and Analysis: HIV Epidemiology Program, NYC DOHMH, 2004


HIV/AIDS testing and prevention

Everyone should know their HIV status. However, an estimated one quarter of New Yorkers living with HIV do not know they are infected, delaying treatment and increasing the risk that they will transmit the disease to others. Less than one fifth of Northeast Queens residents have been tested for HIV in the past year. In addition, half of positive HIV test results (50%) are “late” diagnoses (HIV has already progressed to AIDS) in this community.

The most common way people get HIV is through sexual contact, and having multiple sex partners increases the risk of HIV. Condoms offer protection from HIV when engaging in sexual activities. Less than half of Northeast Queens adults (46%) who had more than 1 sex partner in the past year reported using a condom at their last sexual encounter.


HIV testing

Fewer than 1 in 5 Northeast Queens adults has had an HIV test in the past year



Condom use at last sexual encounter

Less than half of Northeast Queens adults with multiple sex partners used a condom






GOAL 5 Get Help for Depression

Psychological distress

Adults in Northeast Queens are less likely to suffer from serious psychological distress




*Estimate is unstable due to small sample size and should be interpreted with caution.
 Serious psychological distress can be identified in individuals using Kessler's K6 scale, a validated measure consisting of 6 simple questions about mood.
 Percents are age-adjusted.
 Data Source: NYC Community Health Survey 2002-03

Depression is a serious but treatable health condition that frequently goes undiagnosed. Serious psychological distress is associated with depression and other mental illnesses.

In Northeast Queens, residents are less likely to experience serious psychological distress than those in Queens and NYC overall.

Mental illness

Hospitalizations for mental illness are less common in Northeast Queens



Rates are age-adjusted.
 Data Sources: New York State Department of Health Statewide Planning and Research Cooperative System, 1995-2004; U.S. Census 1990 and 2000/ NYC Department of City Planning

Hospitalization rates are one way to look at serious mental illness in a neighborhood. Residents in Northeast Queens have had a fairly constant mental illness hospitalization rate over the past 10 years (excluding alcohol- or drug-related illness).

In 2003-2004, the community's average annual rate of mental illness hospitalizations (381/100,000) was 40% lower than the Queens rate (636/100,000) and less than half the rate in New York City overall (813/100,000).


Understanding hospitalizations and access to health care. Hospitalization data are useful in understanding the burden that certain conditions place on the health care system, but not necessarily in measuring the exact extent of illness in a community. Variations in hospitalization rates may reflect not only differences in rates of illness, but also differences in access to health care. For example, the kinds of health institutions available to residents differ by community, as might the ability of residents to pay for those resources. If a community has a specialized residential institution for a certain type of disease, such as mental illness or stroke, people from outside that neighborhood may come to reside at this institution for care, resulting in an increase in reported hospitalizations for that disease in the community.



GOAL 6 Live Free of Dependence on Alcohol and Drugs

Binge drinking

More than 1 in 10 adults in Northeast Queens engaged in binge drinking in the past month



Percents are age-adjusted.
Data Source: NYC Community Health Survey 2002-03-04

The abuse of alcohol and drugs can lead to many preventable injuries, illnesses, and deaths, including injury in motor-vehicle crashes, liver disease, and violence.

Estimates of binge drinking represent the risk of immediate alcohol-related problems, such as alcohol-poisoning, injury and violence. In Northeast Queens, 14% of adults report engaging in at least one episode of binge drinking (defined as consuming 5 or more drinks on one occasion) in the past month.


Alcohol-related hospitalizations reflect both acute and chronic (e.g., liver disease) consequences of alcohol abuse. The alcohol-related hospitalization rate in Northeast Queens has remained fairly constant in the past decade. In 2003-2004, the average annual alcohol-related hospitalization rate in this community was lower than in Queens and in New York City overall (111/100,000 vs. 235/100,000 in Queens and 439/100,000 in NYC).

The drug-related hospitalization rate in 2003-2004 was also lower (131/100,000) than in Queens (234/100,000) and New York City overall (595/100,000).

The 2002-2004 death rate due to drugs in Northeast Queens was similar to the Queens rate but lower than the rate in NYC overall (2/100,000 vs. 6/100,000 in Queens and 10/100,000 in NYC).

Alcohol-related hospitalizations


The alcohol-related hospitalization rate is lower in Northeast Queens



Rates are age-adjusted.
Data Sources: New York State Department of Health Statewide Planning and Research Cooperative System, 1995-2004; U.S. Census 1990 and 2000/NYC Department of City Planning

Drug-related hospitalizations


The drug-related hospitalization rate is lower in Northeast Queens



Rates are age-adjusted.
Data Sources: New York State Department of Health Statewide Planning and Research Cooperative System, 1995-2004; U.S. Census 1990 and 2000/NYC Department of City Planning

Drug-related deaths

The drug-related death rate in Northeast Queens is lower than the NYC overall rate




Rates are age-adjusted.
Due to small numbers, 3 years of data had to be combined and time trends could not be presented.
Data Sources: Bureau of Vital Statistics, NYC DOHMH, 2002-2004; U.S. Census 2000/NYC Department of City Planning



GOAL 7 Get Checked for Cancer

Cancer screenings

In Northeast Queens, cancer screening is lower than TCNY targets




Percents are age-adjusted.
Data Sources: ¹NYC Community Health Survey 2002 & 2004, ²NYC Community Health Survey 2003-04

Cancer screening can save lives by preventing disease, catching cancer in its early stages and providing opportunities for treatment. TCNY has set specific screening targets for cervical, breast, and colon cancers.

Women in Northeast Queens are getting Pap tests for cervical cancer and mammograms for breast cancer at rates close to the TCNY target of more than 85%. However, only half of adults aged 50 and older in this community have had a colonoscopy in the past 10 years.

Cancer deaths

The death rate due to cancer is lower in Northeast Queens than in NYC overall



Rates are age-adjusted.
Data Sources: Bureau of Vital Statistics, NYC DOHMH, 1995-2004; U.S. Census 1990 and 2000/NYC Department of City Planning

The death rate due to cancer has remained fairly steady in Northeast Queens during the past decade. The 2003-2004 average annual cancer death rate was similar to the Queens rate and more than 20% lower than the NYC overall rate (124/100,000 vs. 137/100,000 in Queens and 161/100,000 in NYC).

Preventing cancer and related deaths. Individuals can reduce their risk of the most common cancers. Never smoking or quitting the habit greatly reduces the risk of lung and other cancers. High colon and breast cancer death rates highlight the importance of getting recommended screenings so treatment can begin early.

The highest cancer-related death rates among men in Northeast Queens are due to lung, blood-related (such as lymphoid), and colon cancers. Among women, breast, lung, and pancreas cancers are the top 3 causes of cancer-related death.

Highest cancer death rates in Northeast Queens (2003-2004)					
MEN			WOMEN		
Type of Cancer	DEATHS / 100,000 PEOPLE		Type of Cancer	DEATHS / 100,000 PEOPLE	
	Northeast Queens	NYC		Northeast Queens	NYC
Lung, trachea, bronchus	33	51	Breast	22	26
Blood-related	20	18	Lung, trachea, bronchus	20	28
Colorectal	16	23	Pancreas	13	9
Prostate	12	25	Blood-related	11	12
Pancreas	10	12	Ovary	10	8


Rates are age-adjusted. Data Sources: Bureau of Vital Statistics, NYC DOHMH, 2003-2004; U.S. Census 2000/NYC Department of City Planning



GOAL 8 Get the Immunizations You Need

Immunizations

Flu shot rates among older adults fall below the TCNY target and pneumococcal (pneumonia) immunizations are even lower



Data Source: NYC Community Health Survey 2002-03-04

Immunizations are not just for kids. Of all the deaths that could have been prevented by vaccination, 99% occur in adults. **Take Care New York** has set a target that more than 80% of adults aged 65 and older will get an annual flu (influenza) shot by 2008. The Northeast Queens flu immunization rate among older adults falls short of the TCNY target by almost 20%.


Immunization rates for pneumonia are lower than those for flu across NYC. Older adults in Northeast Queens are more likely than those in NYC overall to have ever received the pneumococcal vaccine, which protects against one common cause of pneumonia.



GOAL 9 Make Your Home Safe and Healthy

Childhood lead poisoning

Lead poisoning among young children continues to be a problem



Lead poisoning is defined as a blood lead level $\geq 10 \mu\text{g/dL}$.
Data Source and Analysis: Lead Poisoning Prevention Program, NYC DOHMH, 2004


Childhood lead poisoning is a health problem that may be associated with decreased intelligence, learning and behavioral problems, and delayed growth and development. While the number of lead-poisoned children (0-17 years old) in New York City has declined dramatically over the past decade, the Health Department aims to eliminate lead poisoning by preventing children's exposure to lead-based paint and other sources of lead.

In 2004, 10 children in Northeast Queens (5/1,000) were newly identified with lead poisoning (defined as a blood lead level greater than or equal to $10 \mu\text{g/dL}$).

Asthma in Adults and Children

Asthma

Asthma is less common among Northeast Queens adults than adults in NYC overall



*Estimate is unstable due to small sample size and should be interpreted with caution. Percents are age-adjusted.
Data Source: NYC Community Health Survey 2002-03-04


Conditions, or "triggers," in the home environment, such as the presence of second-hand smoke or dust, can cause asthma attacks. Some housing conditions associated with asthma triggers, such as rodent or roach infestation, are more common in some neighborhoods than others.

Adults in Northeast Queens (2%) are less likely to report asthma than those in New York City overall (5%).

Neighborhood asthma hospitalization rates depend in part on the percent of residents who have asthma. However, good medical management of asthma can prevent many asthma-related hospitalizations, and patients can work with health care providers to better control their asthma. Thus, the asthma hospitalization rate can also indicate poor access to health care. Asthma hospitalization rates for adults and children (0-17 years old) in Northeast Queens are lower than in Queens and NYC overall.

Adult asthma hospitalizations


Rates of asthma hospitalization are lower in Northeast Queens



Rates are age-adjusted.
Data Sources: New York State Department of Health Statewide Planning and Research Cooperative System, 1995-2004; U.S. Census 1990 and 2000/NYC Department of City Planning

Child asthma hospitalizations

The child asthma hospitalization rate is lower in Northeast Queens



Data Sources: New York State Department of Health Statewide Planning and Research Cooperative System, 1995-2004; U.S. Census 1990 and 2000/NYC Department of City Planning


TAKE CARE NEW YORK Goal 10 Have a Healthy Baby

The health of babies depends on the health of mothers. Good health care for pregnant women includes high quality prenatal care beginning in the first trimester of pregnancy. The average annual percent of women who received late or no prenatal care in 2003-2004 was lower in Northeast Queens (21%) than in Queens (36%) and NYC overall (28%).

Teenage mothers and their babies face a number of risks. Pregnant teens are more likely to be poor and not complete high school than other teens, and they are more likely to have babies born with low birthweight than older women. The birth rate to teenage mothers has decreased over the past 10 years by more than 60% in Northeast Queens. The average birth rate to teen moms in 2003-2004 in this community (11/1,000) was 85% lower than in NYC overall (75/1,000).

Prenatal care


Northeast Queens mothers are less likely to receive late or no prenatal care



Data Source: Bureau of Vital Statistics, NYC DOHMH, 1995-2004

Teenage mothers

The birth rate to teenage mothers (15-19 years) is lower in Northeast Queens




Data Source: Bureau of Vital Statistics, NYC DOHMH, 1995-2004

Babies born with low birthweight tend to have more health problems than others. In 2003-2004, the average percent of babies born with low birthweight in Northeast Queens was 6% — lower than in Queens (8%) and NYC overall (9%).

Infant mortality (the death of babies in the first year of life) has declined over the past 10 years in NYC. The 2002-2004 rate in Northeast Queens (1/1,000) was lower than the rate in NYC overall (6/1000), meeting the TCNY target.

Low birthweight

The percent of low-birthweight babies is lower in Northeast Queens




Low birthweight is defined as <2,500 grams (5.5 pounds).

Data Source: Bureau of Vital Statistics, NYC DOHMH, 1995-2004

Infant mortality rate (IMR)


The Northeast Queens IMR is lower than in NYC overall



Data Source: Bureau of Vital Statistics, NYC DOHMH, 1993-2004


Neighborhood Health Highlight: Falls among Older Adults

Older adults from Northeast Queens are more likely to have fall-related hospitalizations



Data Sources: New York State Department of Health Statewide Planning and Research Cooperative System, 2003-04; U.S. Census 2000/NYC Department of City Planning

The fall-related hip fracture hospitalization rate in older adults is higher in Northeast Queens



Data Sources: New York State Department of Health Statewide Planning and Research Cooperative System, 2003-04; U.S. Census 2000/NYC Department of City Planning

Every New York City neighborhood has different health concerns. Here we highlight falls among older adults in Northeast Queens.

More than 1 in 3 adults aged 65 and older fall each year, and falls are the leading cause of injury deaths among older adults. Among all NYC neighborhoods, Northeast Queens has the fifth highest rate of hospitalizations from falls — 30% higher than the NYC overall rate (1,517/100,000 vs. 1,148/100,000).

Fractures are one consequence of falls, and hip fractures are the most serious type of fracture. One half of older adults who are hospitalized with a hip fracture cannot return home or live independently after the injury. In Northeast Queens, more than one third (35%) of fall hospitalizations among older adults are for hip fractures. The fall-related hip fracture hospitalization rate in Northeast Queens is nearly 50% higher than the rate in NYC overall (675/100,000 vs. 454/100,000).

TAKING ACTION

Three modifiable risk factors for falls are (1) lower body weakness; (2) problems with balance and walking; and (3) taking 4 or more medications, or any psychoactive medication. Older adults should **exercise regularly** to reduce weakness and increase balance, and **review their medications with a doctor or pharmacist**. Environmental factors also increase risk of falls, and up to one half of all falls happen at home.

For more information on **reducing fall hazards**, go to www.cdc.gov/ncipc/pub-res/toolkit/brochures.htm, or call 311.

Technical notes

Analyses

All analyses were conducted by the Bureau of Epidemiology Services, NYC DOHMH, unless otherwise indicated. All estimates in this report were age standardized to the Year 2000 Standard Population, except for age-specific data and mother-child health indicators. All CHS analyses were done in SUDAAN to account for complex survey design and were weighted to the New York City population according to the U.S. Census 2000.

Data sources

NYS DOH hospitalization data: Includes hospitalizations of NYC residents that occurred anywhere in New York State. Patient zip code was used to classify hospitalizations into 42 neighborhoods. Data from 1995-2003 updated in April 2005; 2004 data updated in July 2005.

Vital Statistics data: Includes births and deaths of NYC residents that occurred within New York City. Data were combined across years to increase statistical stability and average annual rates are presented. In addition, infant mortality rates (IMR) were calculated as 3-year annual averages, and this statistic and others may differ from the presentation in "Summary of Vital Statistics" reports from the Bureau of Vital Statistics, NYC DOHMH.

Community Health Survey data: The NYC Community Health Survey (CHS) is an annual random-digit-dial telephone survey of approximately 10,000 adults in New York City. This profile uses the following datasets from this survey: NYC CHS 2002, NYC CHS 2003, NYC CHS 2004, NYC CHS 2002-03-04, NYC CHS 2002-03, NYC CHS 2002 & 2004, and NYC CHS 2003-04. The combined-year datasets increase statistical power, allowing for more stable analyses at the neighborhood level.

National data and information on falls among older adults were taken from the Centers for Disease Control and Prevention's National Center for Injury Prevention and Control at www/cdc.gov/ncipc/.

Neighborhood Definitions

The 42 NYC neighborhoods are based on the United Hospital Fund definitions of neighborhood, which are specified by zip code. For a complete listing of all 42 neighborhoods and their zip codes, go to nyc.gov/health. The zip codes included in analyses of Northeast Queens are 11361, 11362, 11363, and 11364. Please note that some neighborhoods were combined for statistical purposes in the CHS 2002, CHS 2003 and CHS 2004 datasets to make a total of 33 (2002) or 34 (2003, 2004) neighborhoods. Northeast Queens statistics from these individual-year datasets include data from the neighboring communities of Central Queens, including Fresh Meadows, Hillcrest, and Kew Gardens Hills.

Avoidable Hospitalizations

Data based on Ambulatory Care Sensitive Conditions (called "avoidable hospitalizations" in this report) were calculated using the Agency for Healthcare Research and Quality (AHRQ) classification of inpatient hospitalization data. Conditions in the overall measure include: Diabetes Short-term Complications Admission Rate, Diabetes Long-term Complications Admission Rate, Pediatric Asthma Admission Rate, Chronic Obstructive Pulmonary Disease Admission Rate, Pediatric Gastroenteritis Admission Rate, Hypertension Admission Rate, Congestive Heart Failure Admission Rate, Dehydration Admission Rate, Bacterial Pneumonia Admission Rate, Urinary Tract Infection Admission Rate, Angina without Procedure Admission Rate, Uncontrolled Diabetes Admission Rate, Adult Asthma Admission Rate, and Rate of Lower-extremity Amputation among Patients with Diabetes.

Significance Testing

For all data, 95% confidence limits were calculated for neighborhood, borough, and NYC estimates. If these ranges did not overlap, a significant difference was inferred. This is a conservative measure of statistical difference. This methodology also was used to examine differences between years in neighborhood trend data. *Only robust findings found to be statistically significant are discussed in the text.* In addition, all NYC CHS estimates were evaluated for statistical stability using the relative standard error (RSE). Those estimates with an RSE > .30 are flagged in graphs, "Estimate is unstable due to small sample size and should be interpreted with caution."

TCNY report card

The neighborhood was classified according to where it ranked in comparison to the other 41 NYC neighborhoods with Above Average = rankings 1-10, Average = rankings 11-32, and Below Average = rankings 33-42 where 1 = the best neighborhood score. Rankings were computed by combining (or in some cases, using only one indicator) standardized measures (z-scores) of the following health indicators for each TCNY goal: TCNY#1 - primary care provider, insurance, ED visits; TCNY#2 - current smokers; TCNY#3 - diabetes, obesity, exercise, heart disease hospitalizations, heart disease mortality; TCNY#4 - HIV testing, AIDS mortality; TCNY#5 - serious psychological distress, mental illness hospitalizations; TCNY#6 - binge drinking, alcohol-related hospitalizations, drug-related hospitalizations, drug-related mortality; TCNY#7 - cervical cancer screening, breast cancer screening, colon cancer screening, cancer mortality; TCNY#8 - flu immunization; TCNY#9 - child lead poisoning, adult asthma rates; TCNY#10 - prenatal care, teenage mothers, low birthweight, infant mortality.

Cover Photograph: Cocheron Park, Queens. Photo by Chitra Ramaswamy. Maps by Susan Resnick.

Thank you to all the individuals who contributed to these reports: Sonia Angell, Fatima Ashraf, Birgit Bogler, Shadi Chamany, Louise Cohen, Lorna Davis, Erica Desai, Tamara Dumanovsky, Donna Eisenhower, Jennifer Ellis, Tim Frasca, Stephen Friedman, Renu Garg, Chris Goranson, Leena Gupta, Charon Gwynn, David Hanna, Kelly Henning, Mary Huynh, John Jasek, Qun Jiang, Deborah Kaplan, Adam Karpati, Elizabeth Kilgore, Marty Kim, Vani Kurup, Brooke Levinson, Cortnie Lowe, Jingsong Lu, Xiaowu Lu, Jenna Mandel-Ricci, Thomas Matte, Tina McVeigh, Rachel Miller, Trang Nguyen, Leze Nicaaj, Preeti Pathela, Robyn Philburn, Jane Plapinger, Chitra Ramaswamy, Judy Sackoff, Julia Schillinger, Tejinder Singh, Sally Slavinski, Catherine Stayton, Parisa Tehranifar, William Vaughn, Joshua Volle, Joyce Weinstein, Kellee White, Candace Young, and Regina Zimmerman.

Community Health Profile for Northeast Queens

This report is an updated, expanded second edition of the 2002 Community Health Profile for Northeast Queens.

NEW IN THE SECOND EDITION:

- Take Care New York report card
- Time-trend data on births, hospitalizations, and deaths
- More neighborhood-specific health statistics
- Robust estimates from data through 2004

First and second edition reports on all 42 New York City neighborhoods are available from the New York City Department of Health and Mental Hygiene online or by mail.

Web: nyc.gov/health
Click on "My Community's Health"

Email: profiles@health.nyc.gov

Mail:

Community Health Profiles
New York City Department of Health and Mental Hygiene
Division of Epidemiology
125 Worth Street, Room 315, CN-6
New York, NY 10013

**For more information about health issues
in this report,
please call 311.**

Inside

Community-specific information on . . .

Census Data	2
TCNY Report Card	3
Death Rates	4

Take Care New York

Have a Regular Doctor	5
Be Tobacco Free	6
Keep Your Heart Healthy	6
Know Your HIV Status	8
Get Help for Depression	9
Live Free of Dependence on Alcohol and Drugs	10
Get Checked for Cancer	11
Get the Immunizations You Need	12
Make Your Home Safe and Healthy	12
Have a Healthy Baby	13

Neighborhood Health Highlight	14
-------------------------------	----



New York City Department of Health and Mental Hygiene

Division of Epidemiology, Bureau of Epidemiology Services, 125 Worth Street, CN-6, New York, NY 10013

Michael R. Bloomberg
Mayor

Thomas R. Frieden, MD, MPH
Commissioner of Health and Mental Hygiene

PRST STD
U.S. POSTAGE

PAID
NEW YORK, N.Y.
PERMIT NO. 6174

Copyright©2006

The New York City Department of Health and Mental Hygiene

NYC Community Health Profiles, Second Edition feature information about 42 neighborhoods in New York City.

Suggested citation: Olson EC, Van Wye G, Kerker B, Thorpe L, Frieden TR. Take Care Northeast Queens. NYC Community Health Profiles, Second Edition; 2006; 32(42):1-16.