



NEW YORK CITY DEPARTMENT OF
HEALTH AND MENTAL HYGIENE
Mary T. Bassett, MD, MPH
Commissioner

2015 Alert #8: Increase in Legionnaires' Disease in Queens

- **10 cases of Legionnaires' Disease cases occurred among Queens residents in April-May 2015.**
- **Providers should consider Legionnaires' disease when evaluating patients in Queens presenting with signs of pneumonia.**
 - **Culturing *Legionella* from sputum or bronchoalveolar lavage specimens is the preferred method of diagnosis. Notify your lab to ensure that they use the appropriate culture media to test for *Legionella*.**
 - **Urinary antigen testing is also recommended but should be accompanied by an attempt to isolate *Legionella* by culture.**
- **Report cases promptly to the NYC Health Department and submit all *Legionella* isolates to the Health Department's Public Health Laboratory for serogrouping and molecular typing.**

Please Distribute to All Clinical Staff in Internal Medicine, Pediatrics, Geriatrics, Primary Care, Infectious Diseases, Emergency Medicine, Family Medicine, Laboratory Medicine and Infection Control

May 13, 2015

Dear Colleagues,

In May, an increase in Legionnaires' Disease was noted in the Flushing-Clearview section of Queens. From April to May 2015, 10 cases were diagnosed among Queens residents compared with 4 in May 2014 and 4 in April-May 2013. In the past 5 years, a median of 50 Legionnaires' Disease cases were reported in Queens each year. Within Queens, 9 of the 10 recent case-patients resided in the Flushing-Clearview neighborhoods; in 2014, 7 of 50 total Queens cases were from these neighborhoods. The Health Department is investigating to determine whether or not cases are due to a common exposure. The purpose of this communication is to remind practitioners to test for *Legionella* when evaluating Queens residents who may have community acquired pneumonia.

Caused by the bacteria *Legionella*, Legionnaires' disease is characterized by pneumonia occurring 2-10 days after exposure to an environmental source. *Legionella* is a ubiquitous aquatic organism that grows in warm environments (77°- 108°F). Exposure occurs through inhalation of contaminated aerosols from devices such as cooling towers, whirlpool spas, showers, and faucets, and through aspiration of contaminated water. Person-to-person transmission has not been demonstrated. Groups at high risk include the elderly, cigarette smokers, persons with chronic lung or immunocompromising disease, and persons receiving immunosuppressive drugs. The case-fatality rate has declined substantially since the disease was first described and is estimated to be 8%. Recommended treatment options include macrolide or quinolone antibiotics.

Diagnostic Testing

Culture of the organism from respiratory secretions or tissues is the gold standard for diagnosis. Culture has the added benefit of producing bacterial isolates for molecular typing. With molecular typing of isolates, a patient's isolate(s) can be compared to isolates from environmental sources and other patients to identify a potential source of infection.

Please note the following regarding the diagnosis of legionellosis:

- *Legionella* culture requires specialized media. Please alert your microbiology laboratory that you are considering legionellosis in your patient. The best specimens for culturing *Legionella* are sputum or bronchoalveolar lavage fluid.
- Urine antigen testing is widely available as a rapid method for detecting *Legionella*, but it is most sensitive for *L. pneumophila* serogroup 1. Although *L. pneumophila* accounts for the majority of *Legionella* cases, a negative urine antigen test does not rule out legionellosis due to other species and serotypes. Depending on clinical suspicion, providers should obtain specimens for culture and serology to diagnose legionellosis. Furthermore, urine antigen testing does not yield an isolate for molecular typing to help determine the environmental source.
- Serologic diagnosis requires paired sera collected 3-4 weeks apart to detect a (minimum) four-fold increase in antibody titer to a level > 1:128. A single antibody titer of any level is not diagnostic of legionellosis. For diagnosis, both acute and convalescent sera must be obtained.

There is additional information for clinicians on Legionnaires' disease at the Centers for Disease and Control and Prevention's Legionellosis Resource Site at <http://www.cdc.gov/legionella/index.htm>

Recommendations for Providers

To help the Health Department investigate the increase in increase in Legionnaires' disease in Queens, providers should:

- Maintain a high index of suspicion for legionellosis among all patients with community-acquired pneumonia
- Specifically request both culture and urine antigen testing for *Legionella* when indicated.
- Report all cases to the Health Department by calling the Provider Access Line at 1-866-692-3641.
- Send all cultured isolates to the Health Department's Public Health Laboratory for serotyping and molecular testing. Send isolate and laboratory test request form to:

Public Health Laboratory
455 First Ave, Room 136
New York, NY 10016

If you have any laboratory related questions, please call John Kornblum, Chief of Microbiology, at 212-447-6783.

As always, we appreciate our ongoing collaboration with healthcare providers in New York City to help us address emerging infectious disease concerns.

Sincerely,

Robert Fitzhenry

Robert Fitzhenry, PhD
Bureau of Communicable Disease

Sharon Balter

Sharon Balter, MD
Bureau of Communicable Disease