







Protect your child from these serious diseases that can cause your child to become very ill or hospitalized. Vaccination is your child's best protection against serious illness, or even death, from these diseases.

- | | | |
|-----------------------|-------------------------------|----------------------------|
| Diphtheria | Haemophilus influenzae type b | Hepatitis A |
| Hepatitis B | Influenza | Measles |
| Meningococcal Disease | Mumps | Pertussis (Whooping Cough) |
| Polio | Pneumococcal Disease | Rotavirus |
| Rubella | Tetanus | Varicella (Chickenpox) |

Vaccines:



- ❖ Protect your child against these diseases
 - ❖ Help your child develop immunity without getting sick
 - ❖ Are safe and effective
 - ❖ Are required for your child to attend day care and school

“Start your baby’s shots on time and stay on schedule”

See your child’s doctor, or for information on where your child can be vaccinated, Dial 311, or visit nyc.gov/health

RECOMMENDED PRESCHOOL IMMUNIZATION SCHEDULE, JANUARY 2013

Vaccine	Birth	2 months	4 months	6 months	9 months	12 months	15 months	4 - 6 years
Hepatitis B -Hep B	√	√		√ 6-18 months				
Diphtheria, Tetanus, Pertussis -DTaP		√	√	√			√ 15-18 months	√
Haemophilus influenzae type b -Hib		√	√	√		√ 12-15 months		
Polio -IPV		√	√	√ 6-18 months				√
Pneumococcal -PCV13		√	√	√		√ 12-15 months		
Rotavirus		√ 6-14 weeks	√	√ 6-8 months				
Measles, Mumps, Rubella -MMR						√		√
Varicella -Chicken Pox						√ 12-15 months		√
Hepatitis A -Hep A						√ 12-23 months. Second dose given 6 to 18 months after the 1 st dose		
Influenza -Flu				√	Yearly Two doses given at least four weeks apart for children 6 months to <9 years of age being vaccinated for the first time and for some other children in this age group.			
Meningococcal					√ Children 9 months and older in certain high-risk groups			