


MAP I

TUBERCULOSIS NEW CASE RATES, 1966


Tri-State New York Metropolitan Region

The tuberculosis new case rate in the New York Metropolitan Region is highest in the counties in close proximity to Manhattan. Through the year 1966, New York County (Manhattan) remained the hotbed of tuberculosis incidence. In addition, the Bronx and Kings County (Brooklyn) in New York State and Hudson, Essex and Passaic Counties in New Jersey had tuberculosis problems of very large magnitude.




(Base map through courtesy of the Regional Plan Association)

NEW ACTIVE TUBERCULOSIS CASE RATES
 (PER 100,000 POPULATION)
By Health Center Districts
New York City, 1956 and 1966




NEWLY REPORTED ACTIVE TUBERCULOSIS CASES AND RATES BY ETHNIC GROUP New York City, 1956 and 1966

Percentage of New Active Tuberculosis Cases by Ethnic Group




1956 — 6,137 cases



1966 — 3,663 cases


New Active Tuberculosis Case Rate (Per 100,000 Population) by Ethnic Group



**TUBERCULOSIS PATIENTS UNDER ANTI-BACTERIAL
DRUG TREATMENT IN HEALTH DEPARTMENT CLINICS
AND TUBERCULOSIS PATIENTS
IN HOSPITALS ON DECEMBER 31 OF THE YEAR**

New York City, Alternate Years 1956-1966


Number of Patients



* The census figures shown above for 1966 were recorded as of March 31, 1967.


MORTALITY FROM EIGHT COMMUNICABLE DISEASES New York City, 1965

Number
of Deaths



**MORTALITY RATE
FOR CHRONIC BRONCHITIS* AND EMPHYSEMA
New York City, 1955-1965**

**Death
Rate Per
100,000
Population**



* Deaths from acute bronchitis are not included.