


## TUBERCULOSIS NEW ACTIVE CASE RATES 1967
## TRI-STATE NEW YORK METROPOLITAN REGION

MAP I—The core area (darkest shading) of the entire region continues to show the highest new case rate for tuberculosis, particularly in Manhattan and most of its adjacent counties—Bronx, Kings and Hudson. The new case rate tends to decline as it moves away from the heart of the region.

**LEGEND**

Rate per 100,000 population

| | |
|---|---|
| 0-9 | |
| 10-29 | |
| 30-49 | |
| 50+ | |

(Base map through courtesy of Regional Plan Association)

# NEW ACTIVE TUBERCULOSIS CASE RATES
## BY HEALTH CENTER DISTRICTS
## New York City, 1957 and 1967




PERCENTAGE OF NEW ACTIVE TUBERCULOSIS CASES
BY ETHNIC GROUP
NEW YORK CITY, 1957-1967
Percentage of New Active Cases
70
60
50
40
30
20
10
0
WHITE
NEGRO
PUERTO RICAN
YELLOW
1957
58
59
60
61
62
63
64
65
66
67
YEARS
NOTE: Percentages are based on the total number of new active TB cases for which the ethnic group was reported.


NEW ACTIVE TUBERCULOSIS CASE RATE
BY ETHNIC GROUP
NEW YORK CITY, 1957 and 1967
Rate per 100,000 Population
200
150
100
50
0
1957
1967
All Ethnic Groups
Negro
Puerto Rican
White
ETHNIC GROUP


TUBERCULOSIS PATIENTS UNDER
DRUG TREATMENT IN HEALTH DEPARTMENT CLINICS
AND TUBERCULOSIS PATIENTS IN HOSPITALS
NEW YORK CITY, 1957-1967
Number of Patients
15,000
14,000
13,000
12,000
11,000
10,000
9,000
8,000
7,000
6,000
5,000
4,000
3,000
2,000
1,000
0
COMBINED TOTAL UNDER ACTIVE TREATMENT
CLINIC PATIENTS UNDER DRUG TREATMENT
PATIENTS IN HOSPITAL
1957
58
59
60
61
62
63
64
65
66
67
YEARS

## PERCENTAGE DISTRIBUTION OF DEATHS FROM ALL INFECTIOUS AND PARASITIC DISEASES IN NEW YORK CITY 1966




| TOTAL DEATHS FROM INFECTIOUS AND PARASITIC DISEASES — | 777 |
|---|---|
| Tuberculosis | 537 |
| Septicemia and Pyemia | 66 |
| Syphilis and Its Sequelae | 63 |
| Hepatitis | 19 |
| Acute Infectious Encephalitis | 18 |
| Meningococcal Infections | 12 |
| Measles | 8 |
| All Other Infectious and Parasitic Diseases (including poliomyelitis and diphtheria) | 54 |

Chart based on Section I, Infective and Parasitic Diseases, Numbers 1 through 138; International Classification of Diseases, World Health Organization.


EMPHYSEMA AND CHRONIC BRONCHITIS*
DEATH RATES
NEW YORK CITY, 1955-1966
Death Rate per 100,000 Population
5
4
3
2
1
0
EMPHYSEMA
CHRONIC BRONCHITIS
1955 56 57 58 59 60 61 62 63 64 65 66
YEARS
*Deaths from acute bronchitis are not included