

NUMBER OF DEATHS AND DEATH RATE PER 100,000 POPULATION
FROM SELECTED RESPIRATORY DISEASES IN NEW YORK CITY
1961-1966

Disease and International Code	1961		1962		1963		1964		1965		1966	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
Malignant Neoplasms of Respiratory System Not Specified as Secondary (160-164)	2,767	35.6	2,895	37.2	2,914	37.5	2,949	37.6	2,991	37.6	3,111	38.7
Asthma (241)	234	3.0	213	2.7	227	2.9	190	2.4	197	2.5	201	2.5
Influenza (480-483)	39	0.5	34	0.4	55	0.7	14	0.2	18	0.2	13	0.2
Pneumonia (except newborn) (490-493)	3,466	44.5	3,158	40.6	3,642	46.8	3,223	41.1	3,321	41.7	3,561	44.3
Bronchitis (500-502)	245	3.1	253	3.3	286	3.7	297	3.8	297	3.7	353	4.4
Pneumoconiosis (523-524)	12	0.2	15	0.2	18	0.2	12	0.2	13	0.2	13	0.2
Other Chronic Interstitial Pneumonia (525)	89	1.1	120	1.5	96	1.2	96	1.2	112	1.4	94	1.2
Bronchiectasis (526)	98	1.3	101	1.3	98	1.3	81	1.0	74	0.9	79	1.0
Emphysema Without Bronchitis (527.1)	238	3.1	279	3.6	312	4.0	345	4.4	401	5.0	465	5.8
Other Diseases of the Respiratory System (470-475, 510-522, 527.0, 527.2)	176	2.3	161	2.1	134	1.7	150	1.9	194	2.4	202	2.5

Note: This table includes non-resident deaths, deaths to persons with residence unknown and resident death in institutions. The borough tables do not include these categories and hence the city total will not be equal to the sum of the boroughs.

NUMBER OF DEATHS AND DEATH RATE PER 100,000 POPULATION
FROM SELECTED RESPIRATORY DISEASES IN MANHATTAN
1961-1966

Disease and International Code	1961		1962		1963		1964		1965		1966	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
Malignant Neoplasms of Respiratory System Not Specified as Secondary (160-164)	629	37.0	670	40.6	645	39.1	670	40.4	616	36.2	610	35.4
Asthma (241)	53	3.1	57	3.5	62	3.8	45	2.7	44	2.6	44	2.6
Influenza (480-483)	12	0.7	10	0.6	11	0.7	3	0.2	3	0.2	2	0.1
Pneumonia (except newborn) (490-493)	893	52.6	771	46.7	918	55.6	812	48.9	810	47.6	756	43.8
Bronchitis (500-502)	95	5.6	106	6.4	101	6.1	103	6.2	104	6.1	132	7.7
Pneumoconiosis (523-524)	2	0.1	4	0.2	7	0.4	0	0.0	4	0.2	2	0.1
Other Chronic Interstitial Pneumonia (525)	19	1.1	29	1.8	25	1.5	16	1.0	36	2.1	25	1.4
Bronchiectasis (526)	21	1.2	23	1.4	11	0.7	23	1.4	21	1.2	15	0.9
Emphysema Without Bronchitis (527.1)	44	2.6	70	4.2	76	4.6	82	4.9	81	4.8	91	5.3
Other Diseases of the Respiratory System (470-475, 510-522, 527.0, 527.2)	37	2.2	37	2.2	38	2.3	40	2.4	53	3.1	46	2.7

Note: 1966 is the first year when resident deaths in institutions are not included in the borough totals and hence 1966 data are not altogether comparable with data for the preceding years.

NUMBER OF DEATHS AND DEATH RATE PER 100,000 POPULATION
FROM SELECTED RESPIRATORY DISEASES IN THE BRONX
1961-1966

Disease and International Code	1961		1962		1963		1964		1965		1966	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
Malignant Neoplasms of Respiratory System Not Specified as Secondary (160-164)	458	32.1	493	34.5	529	37.0	533	37.3	537	37.0	482	33.0
Asthma (241)	42	2.9	40	2.8	39	2.7	31	2.2	45	3.1	48	3.3
Influenza (480-483)	6	0.4	4	0.3	7	0.5	2	0.1	4	0.3	0	0.0
Pneumonia (except newborn) (490-493)	613	43.0	533	37.3	693	48.5	541	37.8	588	40.6	483	33.1
Bronchitis (500-502)	34	2.4	33	2.3	60	4.2	56	3.9	50	3.4	45	3.1
Pneumoconiosis (523-524)	1	0.1	3	0.2	1	0.1	3	0.2	4	0.3	4	0.3
Other Chronic Interstitial Pneumonia (525)	18	1.3	27	1.9	23	1.6	28	2.0	39	2.7	23	1.6
Bronchiectasis (526)	17	1.2	18	1.3	19	1.3	13	0.9	16	1.1	18	1.2
Emphysema Without Bronchitis (527.1)	38	2.7	37	2.6	33	2.3	45	3.1	58	4.0	59	4.0
Other Diseases of the Respiratory System (470-475, 510-522, 527.0, 527.2)	17	1.2	13	0.9	20	1.4	29	2.0	29	2.0	20	1.4

Note: 1966 is the first year when resident deaths in institutions are not included in the borough totals and hence 1966 data are not altogether comparable with data for the preceding years.

NUMBER OF DEATHS AND DEATH RATE PER 100,000 POPULATION
FROM SELECTED RESPIRATORY DISEASES IN BROOKLYN
1961-1966

Diseases and International Code	1961		1962		1963		1964		1965		1966	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
Malignant Neoplasms of Respiratory System Not Specified as Secondary (160-164)	799	30.4	826	31.6	797	30.5	774	29.6	826	31.4	867	32.8
Asthma (241)	86	3.3	71	2.7	74	2.8	66	2.5	70	2.7	73	2.8
Influenza (480-483)	14	0.5	11	0.4	19	0.7	2	0.1	6	0.2	1	0.0
Pneumonia (except newborn) (490-493)	1118	42.6	1123	43.0	1157	44.3	1134	43.4	1097	41.7	1200	45.5
Bronchitis (500-502)	54	2.1	48	1.8	56	2.1	70	2.7	76	2.9	80	3.0
Pneumoconiosis (523-524)	5	0.2	4	0.2	3	0.1	2	0.1	3	0.1	4	0.2
Other Chronic Interstitial Pneumonia (525)	26	1.0	33	1.3	20	0.8	25	1.0	14	0.5	19	0.7
Bronchiectasis (526)	33	1.3	36	1.4	37	1.4	25	1.0	16	0.6	22	0.8
Emphysema Without Bronchitis (527.1)	80	3.0	80	3.1	90	3.4	97	3.7	111	4.2	147	5.6
Other Diseases of the Respiratory System (470-475, 510-522, 527.0, 527.2)	45	1.7	34	1.3	28	1.1	38	1.5	60	2.3	69	2.6

Note: 1966 is the first year when resident deaths in institutions are not included in the borough totals and hence 1966 data are not altogether comparable with the data for the preceding years.

NUMBER OF DEATHS AND DEATH RATE PER 100,000 POPULATION
FROM SELECTED RESPIRATORY DISEASES IN QUEENS
1961-1966

Diseases and International Code	1961		1962		1963		1964		1965		1966	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
Malignant Neoplasms of Respiratory System Not Specified as Secondary (160-164)	594	32.8	621	33.4	654	35.2	673	35.5	683	35.4	699	35.8
Asthma (241)	39	2.2	39	2.1	42	2.3	38	2.0	31	1.6	25	1.3
Influenza (480-483)	7	0.4	4	0.2	16	0.9	7	0.4	4	0.2	5	0.3
Pneumonia (except newborn) (490-493)	585	32.3	499	26.8	616	33.1	508	26.8	600	31.1	608	31.1
Bronchitis (500-502)	41	2.3	42	2.3	37	2.0	51	2.7	45	2.3	63	3.2
Pneumoconiosis (523-524)	1	0.1	4	0.2	5	0.3	4	0.2	2	0.1	2	0.1
Other Chronic Interstitial Pneumonia (525)	18	1.0	18	1.0	20	1.1	18	0.9	17	0.9	16	0.8
Bronchiectasis (526)	19	1.1	18	1.0	20	1.1	12	0.6	18	0.9	16	0.8
Emphysema Without Bronchitis (527.1)	52	2.9	61	3.3	87	4.7	82	4.3	106	5.5	113	5.8
Other Diseases of the Respiratory System (470-475, 510-522, 527.0, 527.2)	64	3.5	67	3.6	36	1.9	28	1.5	39	2.0	43	2.2

Note: 1966 is the first year when resident deaths in institutions are not included in the borough totals and hence 1966 data are not altogether comparable with the data for the preceding years.

NUMBER OF DEATHS AND DEATH RATE PER 100,000 POPULATION
FROM SELECTED RESPIRATORY DISEASES IN RICHMOND
1961-1966

Disease and International Code	1961		1962		1963		1964		1965		1966	
	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
Malignant Neoplasms of Respiratory System Not Specified as Secondary (160-164)	68	30.6	82	35.7	71	30.9	71	29.6	102	40.8	93	35.8
Asthma (241)	4	1.8	4	1.7	3	1.3	1	0.4	2	0.8	4	1.5
Influenza (480-483)	0	0.0	3	1.3	1	0.4	0	0.0	1	0.4	2	0.8
Pneumonia (except newborn) (490-493)	103	46.4	103	44.8	98	42.6	91	37.9	89	35.6	73	28.1
Bronchitis (500-502)	8	3.6	4	1.7	7	3.0	4	1.7	8	3.2	4	1.5
Pneumoconiosis (523-524)	1	0.5	0	0.0	0	0.0	2	0.8	0	0.0	0	0.0
Other Chronic Interstitial Pneumonia (525)	3	1.4	4	1.7	1	0.4	1	0.4	3	1.2	2	0.8
Bronchiectasis (526)	6	2.7	1	0.4	7	3.0	4	1.7	2	0.8	2	0.8
Emphysema Without Bronchitis (527.1)	7	3.2	6	2.6	7	3.0	13	5.4	20	8.0	18	6.9
Other Diseases of the Respiratory System (470-475, 510-522, 527.0, 527.2)	4	1.8	1	0.4	4	1.7	4	1.7	2	0.8	9	3.5

Note: 1966 is the first year when resident deaths in institutions are not included in the borough totals and hence 1966 data are not altogether comparable with the data for the preceding years.

DEATH RATES PER 100,000 ESTIMATED POPULATION
For Specified Respiratory Conditions

New York City, 1950-1966

Year	Pneumonia All Forms (490-493)	Lobar Pneumonia (490)	Broncho Pneumonia (491)	Primary Atypical Pneumonia (492, 493)	Influenza (480-483)	Bronchitis (500-502)	Other Chronic Interstitial Pneumonia (525)	Emphysema Without Mention of Bronchitis (527.1)
1950	30.3	11.2	18.4	0.7	0.7	1.5	0.5	0.3
1951	31.0	10.6	19.4	0.9	1.1	1.3	0.6	0.4
1952	32.9	10.7	21.4	0.8	0.5	1.3	0.4	0.3
1953	35.7	10.9	23.6	1.3	1.0	1.4	0.8	0.5
1954	30.1	9.2	19.8	1.1	0.4	1.2	0.9	0.7
1955	34.6	9.9	23.2	1.4	0.6	1.6	1.0	0.8
1956	34.4	10.2	23.4	0.8	0.4	1.8	1.0	0.9
START OF ASIAN INFLUENZA OUTBREAK								
1957	43.9	12.0	30.9	1.0	2.1	2.0	1.0	1.4
1958	45.4	10.1	34.3	1.1	0.8	2.3	1.2	1.8
1959	48.1	9.5	37.2	1.4	0.6	2.4	1.1	2.2
1960	45.7	8.3	36.2	1.2	0.5	2.8	1.4	2.5
1961	44.5	7.7	35.4	1.4	0.5	3.2	1.1	3.1
1962	40.6	7.0	32.4	1.2	0.4	3.3	1.5	3.6
MAJOR INFLUENZA OUT BREAK IN 1963								
1963	46.8	8.2	36.8	1.8	0.7	3.7	1.3	4.0
1964	41.1	7.2	32.1	1.8	0.2	3.8	1.2	4.4
1965	41.7	8.0	31.2	2.5	0.2	3.7	1.4	5.0
1966	44.3	8.8	32.8	2.7	0.2	4.4	1.2	5.8

DEATH RATES PER 100,000 ESTIMATED POPULATION
Pneumonia All Forms (490-493)
By Age Groups-New York City, 1950-1966

Year	AGE GROUPS						Total
	1 Month* to 1 Year	1-4 Years	5-19 Years	20-44 Years	45-64 Years	65 & Over	
1950	196.2	17.7	2.3	7.2	32.3	180.7	30.3
1951	168.1	26.0	2.9	7.3	30.1	186.0	31.0
1952	228.7	30.8	3.2	7.7	30.1	183.3	32.9
1953	224.1	29.8	3.8	8.0	32.3	206.6	35.7
1954	222.0	29.8	3.7	7.1	27.1	155.6	30.1
1955	243.6	26.3	3.9	7.7	27.8	195.6	34.6
1956	236.1	25.7	4.1	7.2	28.6	190.5	34.4
START OF ASIAN INFLUENZA							
1957	234.1	22.5	6.0	11.9	41.5	234.0	43.9
1958	292.2	21.7	3.3	9.8	37.4	258.0	45.4
1959	280.1	21.0	3.7	11.5	37.9	249.7	48.1
1960	276.6	36.2	5.7	10.0	33.1	231.4	45.7
1961	233.4	17.3	3.0	9.1	33.8	247.2	44.5
1962	258.4	18.3	3.9	8.2	30.5	216.8	40.6
MAJOR INFLUENZA OUTBREAK IN 1963							
1963	251.4	17.5	3.3	9.0	35.3	265.6	46.8
1964	222.7	15.2	4.7	8.1	30.4	231.3	41.1
1965	232.3	13.0	4.3	7.5	29.0	244.8	41.7
1966	222.4	14.4	2.8	6.4	31.0	268.5	44.3

* Rates for deaths in the one month to 1 year category are obtained by dividing the Product of 100,000 X the number of deaths in the category by the number of Live Births for the year. Since deaths under 28 days are not included in the count, our rate will be slightly lower than it should actually be.

DEATH RATES PER 100,000 ESTIMATED POPULATION
Lobar Pneumonia-490
By Age Groups-New York City, 1950-1966

Year	AGE GROUPS						Total
	1 Month* to 1 year	1-4 Years	5-19 Years	20-44 Years	45-64 Years	65 & Over	
1950	14.1	2.2	0.5	4.5	17.8	58.5	11.2
1951	7.4	1.1	0.7	4.2	16.6	56.6	10.6
1952	15.2	0.9	0.1	4.0	15.9	59.2	10.7
1953	16.7	2.6	0.5	3.9	15.3	59.5	10.9
1954	13.4	1.9	0.3	3.6	13.0	48.0	9.2
1955	15.2	0.9	0.4	4.2	13.1	53.0	9.9
1956	12.0	1.7	0.2	3.5	14.0	55.8	10.2
START OF ASIAN INFLUENZA							
1957	12.0	1.9	0.7	4.9	17.4	58.3	12.0
1958	15.6	1.3	0.4	3.6	12.4	54.7	10.1
1959	8.9	1.5	0.3	4.2	12.8	42.6	9.5
1960	10.8	3.0	0.8	3.4	9.9	37.8	8.3
1961	6.5	1.5	0.1	2.8	8.5	39.7	7.6
1962	10.3	0.5	0.5	2.2	7.8	38.1	7.0
MAJOR INFLUENZA OUTBREAK IN 1963							
1963	9.5	0.7	0.3	3.2	8.8	44.6	8.2
1964	11.4	0.4	0.3	2.9	7.6	38.7	7.2
1965	11.3	1.1	0.4	2.4	7.1	48.7	8.0
1966	11.1	0.5	0.2	1.9	7.9	55.2	8.8

*Rates for deaths in the one month to 1 year category are obtained by dividing the Product of 100,000 X the number of deaths in the category by the number of Live Births for the year. Since deaths under 28 days are not included in the count, our rate will be slightly lower than it should actually be.

DEATH RATES PER 100,000 ESTIMATED POPULATION
Bronchopneumonia (491)
By Age Groups-New York City, 1950-1966

Year	AGE GROUPS						Total
	1 Month* to 1 Year	1-4 Years	5-19 Years	20-44 Years	45-64 Years	65 & Over	
1950	176.3	15.2	1.7	2.4	13.8	119.7	18.4
1951	155.2	24.5	2.2	2.7	13.2	124.4	19.4
1952	206.1	29.3	3.0	3.4	13.6	120.6	21.4
1953	197.5	25.7	3.0	3.6	15.9	142.0	23.6
1954	195.1	26.9	3.2	3.1	13.0	103.6	19.8
1955	211.5	24.0	3.1	3.0	13.9	137.1	23.2
1956	215.1	23.2	3.7	3.5	14.0	131.6	23.4
START OF ASIAN INFLUENZA							
1957	213.8	20.2	5.1	6.6	23.2	171.2	30.9
1958	267.7	20.0	2.9	5.6	24.0	197.9	34.3
1959	261.7	19.1	3.3	6.7	24.0	201.3	37.2
1960	248.3	32.1	4.8	6.1	22.6	188.7	36.2
1961	225.1	14.7	2.6	5.9	23.5	200.2	35.4
1962	235.2	17.6	3.3	5.3	22.1	173.9	32.4
MAJOR INFLUENZA OUTBREAK IN 1963							
1963	235.7	16.1	2.7	5.1	24.5	212.8	36.8
1964	203.0	13.7	4.2	4.8	21.0	184.1	32.1
1965	203.8	11.4	3.5	4.6	20.1	182.4	31.2
1966	194.3	12.3	2.5	4.0	21.1	198.4	32.8

*Rates for deaths in the one month to one year category are obtained by dividing the Product of 100,000 X the number of deaths in the category by the number of Live Births for the year. Since deaths under 28 days are not included in the count, our rate will be slightly lower than it should actually be.

DEATH RATES PER 100,000 ESTIMATED POPULATION
 Primary Atypical and Other Pneumonia (492,493)
 By Age Groups-New York City, 1950-1966

Year	AGE GROUPS						Total
	1 Month* to 1 Year	1-4 Years	5-19 Years	20-44 Years	45-64 Years	65 & Over	
1950	5.8	0.4	0.1	0.3	0.7	2.5	0.7
1951	5.5	0.4	0.1	0.4	0.8	5.0	0.9
1952	7.3	0.6	0.1	0.3	0.7	3.5	0.8
1953	9.8	1.5	0.3	0.5	1.1	5.1	1.3
1954	13.4	0.9	0.3	0.4	1.0	4.1	1.1
1955	17.0	1.3	0.3	0.6	0.9	5.4	1.4
1956	9.0	0.8	0.1	0.2	0.6	3.2	0.8
START OF ASIAN INFLUENZA							
1957	8.4	0.4	0.2	0.4	0.9	4.5	1.0
1958	9.0	0.4	0.0	0.5	1.0	5.4	1.1
1959	9.5	0.4	0.2	0.6	1.0	6.0	1.4
1960	17.4	1.1	0.2	0.5	0.7	4.7	1.2
1961	1.8	1.1	0.2	0.5	1.8	7.1	1.6
1962	13.3	0.2	0.1	0.7	0.7	4.7	1.2
MAJOR INFLUENZA OUTBREAK IN 1963							
1963	6.0	0.7	0.4	0.7	2.0	8.2	1.8
1964	7.8	1.2	0.2	0.5	1.8	8.5	1.8
1965	17.0	0.5	0.3	0.5	1.8	13.7	2.5
1966	17.0	1.6	0.1	0.5	1.9	14.9	2.7

*Rates for deaths in the one month to 1 year category are obtained by dividing the Product of 100,000 X the number of deaths in the category by the number of Live Births for the year. Since deaths under 28 days are not included in the count, our rate will be slightly lower than it should actually be.

DEATH RATES PER 100,000 ESTIMATED POPULATION
Influenza (480,483)
By Age Groups-New York City, 1950-1966

Year	AGE GROUPS						Total
	1 Month* to 1 Year	1-4 Years	5-19 Years	20-44 Years	45-64 Years	65 & Over	
1950	2.6	0.4	0.4	0.3	0.6	4.1	0.7
1951	1.8	0.9	0.1	0.1	0.9	8.7	1.1
1952	2.4	0.2	0.3	0.5	0.3	2.0	0.5
1953	2.5	0.8	0.2	0.4	0.9	6.0	1.0
1954	0.6	0.4	0.0	0.3	0.2	2.4	0.4
1955	1.8	0.2	0.0	0.6	0.6	2.1	0.6
1956	0.6	0.0	0.2	0.5	0.3	1.2	0.4
START OF ASIAN INFLUENZA							
1957	1.2	1.0	0.8	1.0	3.1	8.0	2.1
1958	0.6	0.4	0.1	0.2	0.9	4.3	0.8
1959	0.6	0.2	0.0	0.1	0.7	3.4	0.6
1960	1.8	0.0	0.0	0.2	0.4	2.5	0.5
1961	1.8	0.2	0.2	0.2	0.2	2.8	0.5
1962	0.6	0.4	0.0	0.0	0.7	2.0	0.4
MAJOR INFLUENZA OUTBREAK IN 1963							
1963	0.0	0.2	0.0	0.0	0.4	5.6	0.7
1964	0.0	0.2	0.0	0.0	0.0	1.1	0.2
1965	1.9	0.0	0.0	0.0	0.2	1.2	0.2
1966	0.0	0.2	0.0	0.0	0.1	1.1	0.2

*Rates for deaths in the one month to 1 year category are obtained by dividing the Product of 100,000 X the number of deaths in the category by the number of Live Births for the year. Since deaths under 28 days are not included in the count, our rate will be slightly lower than it should actually be.

DEATH RATES PER 100,000 ESTIMATED POPULATION
Bronchitis (500-502)
By Age Groups-New York City, 1950-1966

Year	AGE GROUPS						Total
	1 Month* to 1 Year	1-4 Years	5-19 Years	20-44 Years	45-64 Years	65 & Over	
1950	14.1	3.9	0.2	0.2	1.3	6.3	1.5
1951	8.6	3.5	0.2	0.3	1.2	5.5	1.3
1952	4.9	2.2	0.5	0.2	1.7	6.0	1.3
1953	7.4	2.1	0.3	0.1	1.4	8.5	1.4
1954	7.3	1.9	0.1	0.2	1.6	4.7	1.2
1955	15.8	2.3	0.1	0.2	2.0	6.2	1.6
1956	10.8	1.7	0.4	0.5	1.7	8.6	1.8
START OF ASIAN INFLUENZA							
1957	15.6	2.1	0.4	0.4	2.0	8.7	2.0
1958	14.4	2.7	0.1	0.4	2.5	11.2	2.3
1959	8.3	1.9	0.2	0.4	3.2	10.2	2.4
1960	7.2	3.5	0.2	0.2	3.0	14.0	2.7
1961	9.5	1.5	0.2	0.3	3.7	16.7	3.1
1962	13.3	0.9	0.2	0.1	3.8	17.9	3.3
MAJOR INFLUENZA OUTBREAK IN 1963							
1963	9.5	1.1	0.2	0.5	3.9	21.2	3.7
1964	2.4	0.4	0.5	0.5	4.1	23.4	3.8
1965	2.5	0.7	0.2	0.5	4.3	22.5	3.7
1966	6.5	1.1	0.1	0.4	4.7	26.8	4.4

*Rates for deaths in the one month to 1 year category are obtained by dividing the Product of 100,000 X the number of deaths in the category by the number of Live Births for the year. Since deaths under 28 days are not included in the count, our rate will be slightly lower than it should actually be.

DEATH RATES PER 100,000 ESTIMATED POPULATION
Other Chronic Interstitial Pneumonias (526)
By Age Groups-New York City, 1950-1966

Year	AGE GROUPS						Total
	1 Month* to 1 Year	1-4 Years	5-19 Years	20-44 Years	45-64 Years	65 & Over	
1950	0.0	0.0	0.0	0.9	1.3	2.3	0.5
1951	1.2	0.0	0.0	0.6	1.0	3.4	0.6
1952	0.6	0.0	0.0	1.6	0.7	2.0	0.4
1953	0.0	0.2	0.0	3.0	1.5	3.2	0.8
1954	1.8	0.2	0.0	1.7	1.5	4.5	0.9
1955	1.0	0.2	0.0	3.1	1.4	5.4	1.0
1956	0.6	0.2	0.1	2.8	1.6	5.2	1.0
START OF ASIAN INFLUENZA							
1957	0.6	0.2	0.0	1.8	1.5	5.5	1.0
1958	0.6	0.0	0.0	1.4	2.1	6.0	1.2
1959	1.2	0.0	0.0	0.0	1.7	5.9	1.1
1960	0.0	0.0	0.0	0.2	2.1	7.7	1.4
1961	0.0	0.0	0.0	0.2	2.0	5.5	1.1
1962	0.6	0.0	0.0	0.2	2.4	8.0	1.5
MAJOR INFLUENZA OUTBREAK IN 1963							
1963	0.0	0.4	0.0	0.2	1.9	6.5	1.3
1964	1.2	0.2	0.0	0.2	1.9	6.3	1.2
1965	0.6	0.0	0.0	0.3	2.6	6.1	1.4
1966	0.7	0.0	0.0	0.2	1.8	5.9	1.2

*Rates for deaths in the one month to 1 year category are obtained by dividing the Product of 100,000 X the number of deaths in the category by the number of Live Births for the year. Since deaths under 28 days are not included in the count, our rate will be slightly lower than it should actually be.

DEATH RATES PER 100,000 ESTIMATED POPULATION
Emphysema Without Mention of Bronchitis (527.1)
By Age Groups-New York City, 1950-1966

Year	AGE GROUPS						Total
	1 Month* to 1 Year	1-4 Years	5-19 Years	20-44 Years	45-64 Years	65 & Over	
1950	0.0	0.0	0.0	0.9	0.7	1.5	0.3
1951	1.2	0.0	0.1	0.6	0.5	2.3	0.4
1952	0.6	0.0	0.0	0.6	0.8	1.3	0.3
1953	0.6	0.0	0.0	0.6	1.0	3.1	0.5
1954	0.6	0.4	0.0	0.7	1.3	3.2	0.7
1955	0.6	0.0	0.0	1.4	1.4	4.6	0.8
1956	0.6	0.0	0.0	0.3	1.4	5.8	0.9
START OF ASIAN INFLUENZA							
1957	0.0	0.0	0.0	1.8	2.1	8.5	1.4
1958	1.8	0.0	0.0	0.4	3.3	10.1	1.8
1959	0.0	0.2	0.1	0.2	3.6	10.8	2.2
1960	0.0	0.4	0.0	0.2	3.4	15.0	2.5
1961	0.0	0.2	0.1	0.3	4.1	18.1	3.1
1962	0.0	0.0	0.0	0.3	5.1	21.0	3.6
MAJOR INFLUENZA OUTBREAK IN 1963							
1963	1.2	0.0	0.0	0.3	5.6	23.9	4.0
1964	1.8	0.2	0.0	0.6	5.2	27.4	4.4
1965	1.3	0.4	0.0	0.5	6.3	30.9	5.0
1966	3.3	0.0	0.0	0.3	7.7	34.5	5.8

*Rates for deaths in the one month to 1 year category are obtained by dividing the Product of 100,000 X the number of deaths in the category by the number of Live Births for the year. Since deaths under 28 days are not included in the count, our rate will be slightly lower than it should actually be.