NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	1 of 18	
Title: Anthropological Laboratory Analysis	Control No. Revision:	
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 23 March 2021	

1. Policy

The Forensic Anthropology Unit (FAU) responds to consultation requests received from OCME personnel and, on occasion, from external agencies. The FAU shall ensure proper present, examination, and analysis of remains following acceptable practices within a field of Forensic Anthropology.

2. Scop

The roce's soutlin apply to all FAU personnel.

3. Anthropologic Tratory Analyses

There are a plittude consequence ropological examinations that the FAU can perform, depending on the consequences, overall condition of the remains, and the type of analysis requested. The following section briefly summarizes the types of anthropological examinations offered by the FAU.

- Determine if remains are osses thuman, and of medicolegal significance.
- Estimate the Minimum Number of Inc. dals (**NI)
- Estimate the biological profile: set ancest age a death, and stature.
- Describe and interpret pathological conditions and anatomic variants.
- Describe and interpret trauma to include antemory, perimortem and dismemberment.
- Describe and interpret taphonomic changes, including asstmorted damage
- Estimate the Postmortem Interval (PMI).

4. Cleaning Remains and Specimen Removal

4.1 **Cleaning/Macerating Remains:** Remains submitted for anthropological analysis are processed based on their overall condition. The following section summarizes some of the appropriate methods that are available. Information about cleaning or macerating remains shall be recorded on the Basic Case Information Form or an Analytical Notes Form.

Note: All current FAU forms can be found on the Anthropology network drive.

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	2 of 18	
Title: Anthropological Laboratory Analysis	Control No.	Revision:
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 2	23 March 2021

- Skeletal Remains Devoid of Soft Tissue: The remains may be brushed to remove excess dirt/debris. Skeletal remains that are muddy, but are devoid of soft tissue may be wet brushed. Once the remains are sufficiently cleaned, they should be dried in one of the secure Anthropology Labs.
 - Ske tal Material with Soft Tissue: Remains with adherent soft tissue may be disciplified and submerged in warm water with detergents or other appropriate stations remained. Prior to putting the remains in water, the Forensic Anthropologist A) she do remove as much soft tissue as appropriate. The remains should be left to soak at a tested pountil the remains are devoid of soft tissue or until the soft tissue can be more than removed manually. When the remains are sufficiently devoid of soft tissue, hey are more different to be and rinsed off. Any excess soft tissue adhering to the ske ctal elements is removed manually. Once processing is finished the skeletal elements are left in the of the secure Anthropology labs to air dry.
- Cartilaginous Remains: Cartilaginous specimens shall be placed in an evidence container filled with a falling efore attempting to conduct analysis, the specimen should be soaked under ruping weter. See Appendix A: Lab Health and Safety, for the policies and procedure, on handle of working with formalin.
- 4.2 **Specimen Removal:** The medical exact her my request the assistance of the FAU in the removal of specimens. Specimens can be recoved for a variety of reasons including but not limited to, adult or sub-adult age determination rauma, and pathology. See ANTH-001 Evidence Security and Management for the procedure for submitting/receiving evidence from the OCME Evidence/Mortuary Departments.
- 4.3 **Health and Safety:** When cleaning and macerating remains and during specimen removal, FAU personnel are responsible for following the nealth and safety precautions outlined in Appendix A, including but not limited to, we sing the approximately level of personal protective equipment (PPE).

5. Examination Methods

Anthropological consultation requests may require various types of vaminations. The following section outlines typical techniques used by the FAU:

- <u>Macroscopic Examination:</u> Macroscopic examination refers to a visual (gross) examination of remains.
- <u>Microscopic Examination</u>: Microscopic examination refers to a visual examination using magnification provided by a microscope.

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	3 of 18	
Title: Anthropological Laboratory Analysis	Control No.	Revision:
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 2	23 March 2021

- <u>Metric Analysis:</u> Measurements using calipers, osteometric boards, threedimensional coordinate measurement machine/system (digitizer), and tape measures are taken using the appropriate method.
 - <u>Rediographic Examination:</u> Examination of remains using medical imaging chiques.
- Note: AU ersonnel are not responsible for taking radiographs. The OCME Radiography Department is responsible for taking radiographs for casework.
- <u>Éxamine</u> to digital images: Examination of remains through the review of digital potograph

6. Anthropological La oratory Analysis

Based on the compleeness of the remain and/or the examination requested, the analyses outlined below may be performed. When referring to the types of analyses, the FAU shall use appropriate and accepted methors and references. See Appendix B for a current list of the most frequently used remember by the FAU. There is no authoritative body in Forensic Anthropology, however only colished methods shall be used during anthropological laboratory analyses. The FAU does not develop in-house quantitative test procedures nor use non-standard methors for a mination of casework.

- 6.1 **Determining Osseous/Dental versus Nea-osseou Non-dental:** The material shall be examined by macroscopic visual examination, microscopic camination, or evaluation of digital images to assess the presence or absence of transfer or structures that characterize osseous and dental material to include overall size and morph logy, a dmarks, cortical or trabecular structures, density and color. The material may be calculated a radiographic examination or submitted to another unit or agency or other instrument specific procedures.
- 6.2 **Determining Human versus Non-human:** Osseous material shall be examined by macroscopic visual examination, microscopic examination, or through the evaluation of digital images to assess morphology, looking for features or la limarks that are characteristic of human or non-human species based on the examiner attaining and experience in comparative osteology. The osseous material can be compared to information or data from published literature and/or from the FAU comparative non-human skeletal materials.
- 6.3 **Determining Medicolegal Significance:** Determination of medicolegal significance is based on taphonomic and/or contextual indicators. Human remains may be determined not to be of medicolegal significance when they are from historic/prehistoric

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	4 of 18	
Title: Anthropological Laboratory Analysis	Control No. Revision:	
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 2	23 March 2021

archaeological contexts, disturbed cemeteries, or anatomical teaching collections. The FAU analyst shall assess the evidence and document the features and/or context used in making their determination.

- In alyse An analysis of the minimum number of individuals (MNI) shall be completed to check or committeling. If commingling is found, the MNI may be estimated by counting the next repeated to ment or portion of an element. Observations concerning condition, articulation pair-matching, morphology, taphonomy, context, and features of the biological professor that lemains also aid in estimating MNI.
- 6.5 Age at Death estimation relysis of age at death is based on skeletal and dental development is substant aging, and on degenerative skeletal and dental changes for adult aging. Outline below are so le of the most frequently used methods for estimating age at death. The arelyst determines the appropriate method and technique based on the material provided and the condition of the remains.

6.5.1 **Developmental Aging Matho**

- **Dental Development:** Dental cheloppe at and the timing of tooth eruption are utilized for assessing sub-act age cadiographs are taken of the maxilla and mandible to assess unerupted eith and root morphology. The teeth are typically compared to standard denta development tables and figures. Whenever possible, the appropriate methods can table for specific ancestral groups may be used.
- Metric Analysis: Long bone diaphyses develor and grow at predictable rates until the proximal and distal epiphyses fuse to the daphysis. Here are accepted metric methods that utilize measurements of the long bases to accurately estimate skeletal age in immature remains.
- Assessing Epiphyseal Appearance and Union: Appearance and John of epiphyses also occur at predictable rates and is an accumeans of estimating skeletal age in individuals under 25 years. All applicable epiphyses are evaluated to develop a "composite" age estimate. The resulting age estimate may either be reported as terminal (e.g., <18) or as an interval (e.g., 16-20).
- **Medial Clavicles:** The medial clavicles are assessed for the stage of epiphyseal fusion to determine age in teenagers and younger adults.

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	5 of 18	
Title: Anthropological Laboratory Analysis	Control No. Revision:	
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 23 March 2021	

6.5.2 **Degenerative Aging Methods:**

- **Pubic Symphysis:** Observing the degenerative changes to the pubic symphysis is a common method used in estimating age at death for adults. The analyst will document the condition of the symphyseal surface and any effect it may have on the age estimate.
- 4th xibs ternal Ends: Adult aging using the sternal rib end was designed for with the 4th rib sternal end, however when both 4th ribs are unavailable or cannot be sessed the 3rd and 5th-9th ribs can be evaluated instead. In cases when an afternate rib is utilized the analyst will document the rib number in the se note.
- 6.5.3 Additional pethods for age Assessment: Sections 6.5.1 and 6.5.2 summarize the most free ently used technic es for sub-adult and adult aging; however, the analyst may boose to include other techniques or methods in their age assessment that are viewed at the OCME FAU as reputable and accepted by the scientific community (i.e., published in an accepted journal/book).
- Age Constructing the Age interval imation requires an assessment of developmental and degenerate charges from various age indicators. Certain methods are more reliable for particular per ds of life, while others provide a more general indicator of age. The analyst enstructs the age interval based on a composite of the available age indicator The anyst will note which age indicators were used for their assessmen nnal age ate is a matter of expert judgment by synthesizing all available including the rmatio appropriateness of the reference data, familiarity method, condition of the remains, etc.
- Ancestry Estimation: Both cranial and post-cranial non-matric and a etric trans are evaluated for ancestry estimation. Results of ancestry estimation may clude group such as, European (White), African (Black), Hispanic, Asian, Native Aharican, or the esults may be Indeterminate. Outlined below are some of the most frequently, sed modes for estimating ancestry; however, the analyst determines the appropriate method and technique based on the material provided and the condition of the remains.
 - 6.6.1 **Non-Metric Assessment of Ancestry**: Non-metric characteristics of the cranium, mandible, and dentition are used when assessing ancestry.
 - 6.6.2 **Metric Assessment of Ancestry**: Statistical software programs, such as Fordisc, are used for metric assessment of ancestry. Bone measurements are taken using an

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	6 of 18	
Title: Anthropological Laboratory Analysis	Control No. Revision:	
	ANTH-002	3
		22.4
Approved by: Forensic Anthropology Director	Effective Date: 23 March 2021	

approved reference (e.g., UT Data Collection Procedures, Howell's definitions). The measurements used for the statistical analysis are up to the discretion of the user. All steps in the statistical software analysis are retained in the associated log files which are kept with the case file (either hardcopy or electronic copy). Any assurement involving trauma, damage, pathologic condition, or anatomical triant is included or excluded at the discretion of the analyst and a comment is nade in analytical notes.

- 6.7 **Sex Estimaton:** Sex estimation is performed by standard non-metric and/or metric assessment procedures that examine sexually dimorphic characteristics of the skeleton. Provided below in an address for sex estimation, however the analyst determines the appropriate bethod a straigue based on the material provided and the condition of the remains.
 - 6.7.1 **Non-metric bethods**: Morphological features of the pelvis and skull are typically used to estimate sex. In addition, the skeletal elements present may be evaluated for overall robusts.
 - 6.7.1 **Metric Analysis**: Estivation of some determined using measurements of the cranial, mandibular, and postcranial elements.
- 6.8 **Stature Estimation**: Stature can be estimated using rathematical methods (e.g., Fordisc) or anatomical methods (e.g., Fully method). Take measurements as described for the method and select the appropriate demographic can gories training, report the 90% prediction interval. Stature may be reported in cerum 1.8, inches a feet and inches.
- 6.9 **Dental examination**: Examine and chart the dentition for an intory process. Dental analyses regarding age, sex, ancestry, or trauma will be provided to the relevant sections. Chart the dentition using the Universal Numbering System and document the cowing:
 - Antemortem tooth loss/agenesis.
 - Postmortem tooth loss.
 - All restorations.

Note: All dental radiographs are taken by the OCME Radiology Department. In most cases, an OCME forensic odontologist will also examine and chart the dentition for identification purposes.

- 6.10 **Pathological Conditions**: Examine and document characteristics of pathological change. At a minimum the following should be documented (when applicable):
 - Affected elements and approximate location.
 - Presence of bone remodeling and extent of healing.

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	7 of 18	
Title: Anthropological Laboratory Analysis	Control No.	Revision:
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 2	23 March 2021

- Presence of accompanying features.
- 6.11 **Anatomical Variants**: Examine the remains for anatomical variants such as abnormal development or notable variations of normal human skeletal anatomy. Describe the anatomy and location.
- 6.12 The val Analysis: Remains are examined for trauma in cases where a full skeletal analysis is presented or on specimens removed from autopsy at the request of the medical examiner. The una analysis involves examining the remains for antemortem, perimore in, an dismember trauma.

The following eneral in should be recorded when describing and interpreting trauma:

- A determination of the training as antemortem, perimortem, or dismemberment, if possible.
- The location of the trauma.
- If antemortem, a description of any healing, signs of medical intervention, and if possible a relative age of njury.
- Description of the type of traum. possible (e.g., blunt, sharp, high velocity projectile).
- Notes on whether a reconstruction of the spatimen is required to perform the analysis.
- Notes on relevant postmortem damage.
- Notes on any relevant pathological conditions the may be associated with the trauma.
- 6.12.1 **Blunt Force Trauma:** The following is specify information that should be recorded for blunt force trauma analysis:
 - Description of the fracture(s) including anator cal location
 - If possible, notes on the direction of force, specific apact site and fracture patterns.
 - If possible, a determination of tool class characteristic minimum amber of impacts, and sequence of impacts.
- 6.12.2 **Sharp Force Trauma:** The following is specific information that should be recorded for sharp force trauma analysis:
 - Descriptions and/or drawing of the location of the defect.
 - Any relevant measurements.
 - Descriptions of specific characteristics of the defect (e.g., incomplete cuts, kerf wall, kerf floor, striation patterns).

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	8 of 18	
Title: Anthropological Laboratory Analysis	Control No. Revision:	
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 2	23 March 2021

- Notes on the progression of the weapon through osseous and cartilaginous structures, when applicable.
- When appropriate, casts of the tool mark (see section 6.12.4).
- If possible, determine tool class characteristics, minimum number of impacts, and sequence of impacts.
- me circumstances, it may be necessary to expose the cut surface (kerf (alls) by cutting the cartilage or bone to open up the defect for floor note: hen this occurs the newly cut surface must be noted in the ation.` o it can be identified as an examination modification.
- 6.12.3 **High** Y locity Trauma: The following is specific information that ed for high velocity projectile trauma analysis: should
 - tions and/ordrawing of the overall shape and anatomical location lefect (wound) with associated fractures.

 In pents of the see of the defect(s).
 - Measurements of the
 - c characteristics of the defect and notes on the Descriptions of s trajectory, if po
 - Statements of the minimum amber defects and sequence of defects, if possible.
- 6.12.4 Tool Mark Casting: FAU analyst May cho se to create cast impressions to aid in examination of tool mark characteristics. polyviny loxane (e.g., Accutrans) or similar casting material is used to creat tool and capaced in bags labeled with the unique of camber and All casts should be iption of the cast. Tool mark casts created during analysis are consider and the FAU evide shall follow the policies and procedures docume Evidence Security and Management (see ANTH-001).
- 6.13 Postmortem Interval and Taphonomic Changes: Examine to remain anv information from the scene that may aid in the assessment of the pos ortem interv and d if possi taphonomic processes. Describe the condition of the remains, the probable interval between death and discovery.
- 6.14 Postmortem Damage: Postmortem damage refers to any damage to the remains after death and can sometimes be misconstrued as perimortem trauma. The following is specific information that should be recorded for postmortem damage, when possible:
 - Documentation of the location of the damage in the analytical notes or, if necessary, in a diagram.
 - Description of the extent, pattern, and possible cause of the damage.

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	9 of 18	
Title: Anthropological Laboratory Analysis	Control No. Revision:	
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 2	23 March 2021

- Notes on taphonomic changes to the remains (e.g., color changes, animal activity, water damage).
- Notes on damage resulting from standard autopsy protocol, which are included within the postmortem damage description, when applicable.

7. Prifyi g New Methods

ed methods shall go through a verification process prior to being used U shall verify that all analysts are competent to use the new method analysi independently perform the method on the same sample(s) and erification is considered complete and the new method can be men all the analysts' results are in agreement. If there is a used on casev as a group, the FAU shall review all results, as well as disagreement b ween re sure all analysts understand how to appropriately use the procedures or the nethod to the method. Contin ed testing the method will occur until all analysts are in agreement.

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	10 of 18	
Title: Anthropological Laboratory Analysis	Control No. Revision:	
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 23 March 2021	

Appendix A. Health and Safety

Policy and Scope

FAU personnel interns and visiting scientists are responsible for following the health and safety policies and put edures outlined by the OCME Health and Safety Department and the safety present as provided in this appendix.

OCME He in an San w Department: The OCME Health and Safety Department is responsible or the salth and afety of all OCME employees. FAU personnel are responsible for following the broth and safety policies and procedures that apply to their duties. OCME health and safety policies and procedure are located on the OCME intranet under Libraries/Health and Safety.

FAU Safety Officer: The AU Quality Assurance (QA) Specialist is the designated Safety Officer for the unit (see the AU organization chart in QM-001: Personnel). The QA Specialist is the primary liaison between he OCME Here I and Safety Department and the FAU. The QA Specialist is responsible for making sure to FAU follows the OCME health and safety policies and procedures as well as the policies and procedures applicitly stated in this appendix. The QA Specialist is also responsible for chemical hygien as a safety issues.

Note: The duties and tasks associated with anntaining heart and safety compliance can be divided among FAU employees.

FAU Personnel: It is the responsibility of FAU personnel of complete the and enforce the health and safety standards created by the OCME Health and after pepartment and outlined in this Appendix.

FAU Laboratory Safety Precautions:

Personal Protective Equipment (PPE): FAU personnel, interand visiting tists are responsible for wearing the appropriate level of PPE required when v rking in the ology Laboratories. The appropriate PPE may vary depending on the task at and PE may in ude, but is not limited to lab coats, scrubs, disposable aprons, disposable gloss, shoe cover eve protection, and respiratory protection. In addition to PPE, FAU personnel shall follow the JCME Laboratory Dress Code policy. Closed toed shoes should be worn when enterarea, working with hazardous materials, or working with sharp instruments. The OCME Laboratory Dress Code policy is located on the OCME intranet under Libraries/Health and Safety.

PPE in Autopsy Suites: At minimum, FAU personnel shall wear appropriate lab attire and an N95 or equivalent face mask upon entering the autopsy suites when autopsies are being performed. The minimum PPE required when working in the autopsy suites may include but is not limited to, a lab coat, disposable apron, disposable gloves, shoe covers,

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	11 of 18	
Title: Anthropological Laboratory Analysis	Control No. Revision:	
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 2	23 March 2021

and an N95 or equivalent face mask. All PPE, except lab coats should promptly be removed and discarded when exiting the autopsy suite or morgue area. Disposable PPE should be discarded in the designated red biohazard bins.

PPE of Vorking in Anthropology Laboratories: When working in the Anthropology aboratories, FAU personnel shall wear PPE appropriate to the task. The type of PPE who yellow the personnel shall wear PPE appropriate to the task. The type of PPE who yellow depending on the task(s) being performed (e.g., maceration may require additional PPE than is not necessary when conducting skeletal analyses). Disposable gloves work to handle processed specimens (i.e., dry bones) can be discarded in the lab garbage ans.

Note: Let mac ation problem before handling or use the oven mitts when handling heated pots.

Sharps Safety: FAU personnel using sharp instruments (e.g., scalpels and bone saws) with possible exposure to body flute. The regarded to wear cut gloves underneath their disposable gloves.

Formaldehyde/Formalin Exposure: When yoking the formalin fixed specimens FAU personnel shall follow the policies described it the NY OCM Chemical Safety Plan. The most up-to-date version of the Chemical Safety Planes local on the OCME intranet under Libraries/Health and Safety.

When working with formalin fixed specimens the foll yirr procedure ould be adhered to whenever possible:

- Prior to examination the specimen(s) should be soaked under an ing wat
- At minimum wear a lab coat or disposable apron and disposable nin le gloves.
- To avoid inhalation of formaldehyde fumes all containers alled with forman pould be closed at all times, except when removing or returning specime (s) to the ontainer

Chemical Hygiene: Chemical hygiene refers to working with and handing chemical in a hygienic or clean manner. FAU personnel are responsible for handling an using remicals properly from initial receipt to final disposal.

All hazardous chemical containers shall be initialed and dated when received and first removed from their shipping containers. A proper notation in the "Chemical Inventory Form" shall be completed for each chemical received. The QA Specialist shall review this form as part of his/her annual audit or when deemed necessary. Additionally, the QA Specialist shall make sure that Safety Data Sheets (SDS) for hazardous chemicals are accessible to all FAU personnel.

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	12 of 18	
Title: Anthropological Laboratory Analysis	Control No. Revision:	
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 2	23 March 2021

Handling Hazardous Chemicals: FAU personnel shall:

- Wear proper PPE to avoid skin/eye contact with hazardous chemicals.
- Wash their hands after handling chemicals.
- But and it with the SDSs of any hazardous chemicals used. These sheets are available line; and in the FAU Chemical Records Binder.
- Pur vardous a micals back into the chemical storage cabinet after use (see Chemical Storage).
- Discusse of Anardous Cemicals properly (see Chemical Disposal).

Chemical Storage: He areads chemicals shall be stored in the FAU chemical storage cabinet. The FAU chemical storage cabinet are used in the Anthropology Laboratory located on the 4th floor (room 424).

Chemical Disposal: Expire or deteriorated of micals or chemicals no longer utilized shall be disposed of properly. The Come Health of Safety Department should be consulted prior to chemical waste disposal and an OcME Comical Waste Removal Tracking Sheet shall be filled out and forwarded to Health and Safety from the lisposal.

Chemical Records: Records related to FAU comicals add as the Chemical Inventory Form, OCME Chemical Waste Removal Tracking Seet, and affety at a Sheets shall be maintained by the QA Specialist. A chemical inventory shall be completed by the QA Specialist during the annual audit.

Incident Reporting: Any accident with injury shall be as wed first at the reported to the Forensic Anthropology Director (Director) and the OCME Health and affety a partment. When reporting an incident, the Director and injured personnel shall follow as steps at lined on the OCME Injury or Illness at Workflow chart. The most current version of the Injury and Illness at Workflow chart is located on the OCME intranet under Libraries/Health and Safety

Housekeeping: Each FAU employee is responsible for the cleanliness of his ter workspal and jointly responsible for the Anthropology lab/office spaces.

The following procedures apply to the housekeeping standards of the laboratory:

- The Anthropology labs shall be kept clean and orderly. Any spills or messes shall be cleaned immediately.
- All lab equipment shall be kept in their assigned storage areas, except when in use.
- All chemical and biological waste shall be disposed of properly.
- Pathways, doorways, fire-extinguishing equipment and any other emergency equipment shall remain unobstructed.

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	13 of 18	
Title: Anthropological Laboratory Analysis	Control No.	Revision:
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 2	23 March 2021

Appendix B. Current list of References

Al Qahtani, S.J., Hector, M.P., and Liversidge, H.M.

Brief Communication: The London Atlas of Human Tooth Development d Eruption. American Journal of Physical Anthropology 142(3):481-490.

Aufo neide C. d C. Rodríguez-Martín

199 The Carbridge Encyclopedia of Human Paleopathology. Cambridge: mbh. ve University Press.

Barnes, E.

2013 Itlas developmental field anomalies of the human skeleton: A paleopatholo perspective Hoboken, NJ: John Wiley and Sons.

Bass, W.M.

2005 Human Os. Logy. umbia: Missouri Archaeological Society.

Birkby, J.E., Fenton, T.W., and B.E., And and A.

2008 Identifying southwest Historics yang Nonmetric traits and the cultural profile. Journal of Forensic Sciences 53 (29-3).

Brooks, S., and Suchey, J.M.

1990 Skeletal age determination based upon the optable; a comparison of the Acsadi-Nemeskeri and Suchey-Brooks me took atuman Fortuin 5(3):227-238.

Buikstra, J.E., and D.H. Ubelaker, eds.

1994 Standards for Data Collection from Human skeleta. Semains, Proceedings of a Seminar at the Field Museum of Natural History (Arkansas Archaeola et al. Survey Research Series No. 44). Fayetteville: Arkansa Archaeola et al. Survey.

Browner, B.D., Jupiter, J.B., Levine, A.M., and P.G. Trafton, eds.

2003 Skeletal Trauma: Basic Science, Management, at Recondition.
Philadelphia: Saunders.

Cramer, K.E., and N.E. Green, eds.

2003 Skeletal Trauma in Children. Philadelphia: Saunders.

Crowder, C., C.W. Rainwater, and J.S. Fridie

2013 Microscopic analysis of sharp force trauma in bone and cartilage: A validation study. Journal of Forensic Sciences 58(5):1119-1126.

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	14 of 18	
Title: Anthropological Laboratory Analysis	Control No.	Revision:
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 2	23 March 2021

Edgar, H.J.H.

2005 Prediction of race using characteristics of dental morphology. Journal of Exercise Sciences 50:269-273.

zekas I., and Kósa, K.

1978 Fensic Fetal Osteology. Budapest: Akadémiai Kiadó Publishers

Gal way,

74 Token Jones. Springfield, Charles C. Thomas.

Gill, G.W., and thine, S

1990 keletz attribution of race: Methods for forensic anthropology. Maxwell Museum Antropological I pers no. 4. Albuquerque: University of New Mexico.

Haglund, W.D., and N.H. Sorg, eds.

2002 Advances Sorens' Caphonomy. Boca Raton: CRC Press.

Humphrey, J.A. and D.L. Hucknson

2001 Macroscopic characterist of locking trauma. Journal of Forensic Sciences 46(2): 228-233.

Hartnett, K.M.

2010 Analysis of Age-at-Death Estimation Using and from a New, Modern Autopsy Sample-Part I: Pubic Bone. Journal of Forensia Science 55(5):1145-1151.

2010 Analysis of age-at-death estimation using data om a new, modern autopsy sample—part II: sternal end of the forth rib. Sournal of Forensic Sciences 55(5):1152-1156.

Hefner, J.T.

2009 Cranial nonmetric variation and estimating ancestry. Turnal of Grensic Sciences 54:985-995.

Hefner, J.T., Ousley, S.D,

2014 Statistical classification methods for estimating ancestry using morphoscopic traits. Journal of Forensic Sciences 59(4):883-890.

Hurst, C.V.

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	15 of 18	
Title: Anthropological Laboratory Analysis	Control No.	Revision:
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 23 March 2021	

2012 Morphoscopic trait expressions used to identify Hispanic ancestry. Journal of Forensic Sciences 57(4):859-865.

İşcan, M.Y., and R.K. Wright.

1 3 Casts of age phases from the sternal end of the rib for white males and males. Fort Collins, Colorado: France Casting.

İşcar M.Y., V. an, R., and R.K. Wright

1984 Age est nation from the rib by phase analysis: white females. Journal of Lens, Sciences 30(3):853-863.

İşcan, M.Y., Lon, S.R., Wright

1984 Letam phosis at the sternal rib end: a new method to estimate age at death in white males. American Journal of Physical Anthropology 65(2):147-156.

Jantz, R.L., and S.D. usley

2005 FORDISC composite discriminant functions. Version 3.0.: The University of Tennesce, Legaville

Katz, D. and Suchey, J.M.

1986 Age determination of male s pulls. American Journal of Physical Anthropology 69:427-435.

Kenyhercz, Michael W., et al.

2017 The Optimized Summed Scored A trib as Method or the Classification of U.S. Blacks and Whites: A Validation Study. Joynal of Sprensic Sciences 62(10: 174-180.

Klales, A.R.

2018 MorphoPASSE: the Morphological Pelvis of Skull ox Estimation Database. Topeka, KS: Washburn University.

Klales, A.R and S.J. cole

2018 MorphoPASSE: the Morphological Pelvis and Skull Stimation Database Manual. Topeka, KS: Washburn University.

Klales, A.R., S.D. Ousley, and J.M. Vollner

2012 A revised method of sexing the human innominate using Phenice's nonmetric traits and statistical methods. American Journal of Physical Anthropology 149(1):104-114.

Kleinman, P.K., ed.

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	16 of 18	
Title: Anthropological Laboratory Analysis	Control No.	Revision:
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 23 March 2021	

2015 Diagnostic Imaging of Child Abuse. 3rd edition. Cambridge: Cambridge University Press.

Komar, A.

Decay rates in a cold climate region: a review of cases involving advanced composition from the medical examiner's office in Edmonton, Alberta. Journal of Foren Sciences 43(1):57-61.

Lan ey, N. Jantz, M., Ousley, S.D., Jantz, R.L., and G. Milner

Anthrop Collection Procedures for Forensic Skeletal Material. Department of Anthrop Collection Procedures for Forensic Skeletal Material. Department of Anthrop Collection Procedures for Forensic Skeletal Material.

Langley-Shirle N., 2 A.L. Jan

2010 A Ba sian approach to age estimation in modern Americans from the clavicle. Journal of Forensic Sciences 55(3):571-583.

Love, J.C., S.M. Derrick, J. W. Jema

2011 Skeletal Atlas of Alla buse. Ww York: Humana Press.

Mann, R.W., Jantz, R.L., Bass, W., and P. Wille

1991 Maxillary suture obliter from: a sual rethod for estimating skeletal age. Journal of Forensic Sciences 36(3): 7 2-791.

Meindl, R.S., and C.O., Lovejoy

1985 Ectocranial suture closure: a rey editathod force determination of skeletal age at death based on the lateral anterior sures. As rican Journal of Physical Anthropology 68(1): 57-66.

Mincer, H.H., Harris, E.F., and Berryman, H.E.

1993 The A.B.F.O. study of third molar developmen and its use as estimator of chronological age [published erratum appears in Journal Fore at Sciences 1993 Nov; 38(6):1524]. Journal of Forensic Sciences 38(2):379-36

Ortner, D.J.

2003 Identification of pathological conditions in human skeletal remains. San Diego: Academic Press.

Phenice, T.W.

1969 A newly developed visual method of sexing the os pubis. American Journal of Physical Anthropology 30(2):297-301.

Pokines, J., and Symes, S.A.

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	17 of 18	
Title: Anthropological Laboratory Analysis	Control No.	Revision:
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 23 March 2021	

2013 Manual of Forensic Taphonomy. Boca Raton: CRC Press.

Reichs, K.J.

1998 Forensic Osteology: Advances in the Identification of Human Remains. Singfield, Charles C. Thomas.

Roe L.J., and R., Yochum, eds.

2005 Associals of Skeletal Radiology. 3rd edition. Philadelphia, Lippincott Williams & Wokins.

Scheuer, L. and Black

2004 he Juranie Skele n. San Diego, Elsevier Academic Press.

Shirley, N.R., Jantz L.

2011 Spheno-scipital Synch drosis Fusion in Modern Americans

Symes, S.A.

1992 Morphology & Saw March Human Bone: Identification of Class Characteristics. Dissertation, Anthopology University of Tennessee.

Ubelaker, D.H., and B.J. Adams

1995 Differentiation of perimor in and perimor trauma using taphonomic indicators. Journal of Forensic Sciences 40(4): 509-51

Walker, P.L.

2008 Sexing skull using discriminant analysis a sually sessed traits. American Journal of Physical Anthropology Journal 136(1), 9-50.

Yoder, C., Ubelaker, B.A., and Powell, J.F.

2001 Examination of Variation in Sternal Rib End Morpholo Relevant Age assessment. Journal of Forensic Sciences 46(2):223-227.

Revision History

REV.	DATE	SUMMARY OF CHANGES
0	26 January 2018	New document.
1	18 October 2018	Document Control No. changed from ANTH-003 to ANTH-002.

NYC – OFFICE OF CHIEF MEDICAL EXAMINER	Page:	
Quality Management	18 of 18	
Title: Anthropological Laboratory Analysis	Control No.	Revision:
	ANTH-002	3
Approved by: Forensic Anthropology Director	Effective Date: 2	23 March 2021

		6.0- Added the following statements: "There is no authoritative body in Forensic Anthropology, however only validated and published methods shall be used during anthropological laboratory analyses. The FAU does not develop in-house quantitative test procedures nor use non-standard methods for examination of casework. Created Section 7. Verifying New Methods: Any newly validated, published methods shall go through a verification process prior to being used on casework. FAU analysts shall perform verification of a new method by cticing/testing the method on sample(s) and comparing the autome to the results from one of our already apprayed analytical methods.
2	27 February 2020	Appendix /* Edited the following statement in the PPE in Autopsy Suite section: "A PPE except be coats should promptly be removed and discard depen exiting the autopsy suite or morgue area. Dispose the PPE mould be discarded in the designated red biobased bip. Added the following statement to the PPE for working in Anthropological aboratories section: "Disposable gloves from to be the processed specimens (i.e., dry bones) can be discarded in a lab garbage bins."
3	23 March 2021	Removed the following sentence for clause 4, "all specimens removed shall be submitted to a Evidence Department. Removed the word "validated from clause 6. And In Appendix A, under the Forman shyde sections of the bullet point was removed. In Appendix B, several references were added to the citation list and the Howell 1973 citation was removed.